



PO Box 201800 • 1201 11th Avenue • Helena, MT 59620-1800 • fax 406.444.0266 • tel 406.444.5363 • <https://mtnhp.org>

October 17, 2023

Tamara Ross
IMEG Corp.
1817 South Ave West
Missoula, MT 59801

Dear Tamara Ross,

Thank you for your request for Natural Heritage information for the Gooden Curtis Subdivision, within Section 20, T13 N R19 W, in Missoula County. Included with this letter is an Environmental Summary report PDF and a companion Excel workbook summarizing information managed in the Montana Natural Heritage Program's (MTNHP) databases for: (1) species occurrences; (2) other observed species without Species Occurrences; (3) other species potentially present based on their range, presence of associated habitats, or predictive distribution model output if available; (4) structured surveys (organized efforts following a protocol capable of detecting one or more species); (5) land cover mapped as ecological systems; (6) wetland and riparian mapping; (7) land management categories; and (8) biological reports associated with plant and animal observations. The PDF report contains introductory materials and limitations associated with the use of each of these data types, a list of additional information resources, data use terms and conditions, and suggested contacts. The Excel workbook contains worksheets for each data type that can be easily sorted to summarize particular information needs. In addition to these materials, we have included a compilation of one page snapshots containing general description, habitat, spatial and temporal distribution, and conservation status information for each species listed in the species occurrence, other observed species, and other potential species sections of the Environmental Summary report. These three field guide compilations are excerpted from the full accounts found on the Montana Field Guide <https://fieldguide.mt.gov> for general reference use and, if desired, as appendices to environmental review documents.


Please keep in mind the following when using and interpreting the enclosed information:

- (1) This information is intended for distribution or use only within your department, agency, or business. Please see the Data Use Terms and Conditions in the Environmental Summary report PDF for additional guidelines.

- (2) Our minimum search area for standard information requests consists of the requested area buffered by an additional mile in order to capture records that may be immediately adjacent to the requested area. Please let us know if a buffer greater than 1 mile would be of use to your efforts.
- (3) Additional information on animal, plant, and lichen species and ecological systems in Montana is available on the Montana Field Guide at <https://fieldguide.mt.gov/>
- (4) In addition to the information you receive from us, we encourage you to contact state, federal, and tribal resource management agencies in the area where your project is located (see Environmental Summary report PDF).

I hope the enclosed information is helpful to you. Please feel free to contact me at the phone or email address below if you have any questions, require additional information, or have suggestions for how we could improve our information resources.

Sincerely,

A handwritten signature in blue ink that reads "Bryce A. Maxell".


Bryce A. Maxell
Montana Natural Heritage Program
(406) 444-3989
bmaxell@mt.gov



Montana SOC Occurrences Report

Report generated 10/17/2023 11:18:58 AM

SOC Occurrences for Birds = Bald Eagle



Birds - Bald Eagle (*Haliaeetus leucocephalus*)

SO Count: 9

Obs Count: 119

Earliest Obs: 2004

Recent Obs: 2022

Special Status Species
Native Species

Global Rank: G5

State Rank: S4

Agency Status

USFWS: BGEPA; MBTA

 USFS: Sensitive - Known in
 Forests (BD, BRT, KOOT,
 LOLO)

BLM: SENSITIVE

FWP SWAP:

PIF: 2

Delineation Criteria

Confirmed nesting area buffered by a minimum distance of 2,000 meters in order to be conservative about encompassing the breeding territory and area commonly used for renesting. Only nesting observations with a locational uncertainty of 1,000 meters or less will be used to delineate a nesting area.

Last Updated

Sep 05, 2023

<input type="checkbox"/> SO ID: 51159192	Acres: 3,095	Obs Count: 3	Earliest Obs: 2006	Recent Obs: 2021
<input type="checkbox"/> SO ID: 51159296	Acres: 3,105	Obs Count: 4	Earliest Obs: 2005	Recent Obs: 2006
<input type="checkbox"/> SO ID: 51159526	Acres: 3,095	Obs Count: 7	Earliest Obs: 2016	Recent Obs: 2022
<input type="checkbox"/> SO ID: 51159635	Acres: 3,105	Obs Count: 18	Earliest Obs: 2006	Recent Obs: 2017
<input type="checkbox"/> SO ID: 51160348	Acres: 3,105	Obs Count: 1	Earliest Obs: 2007	Recent Obs: 2007
<input type="checkbox"/> SO ID: 51160351	Acres: 3,095	Obs Count: 62	Earliest Obs: 2008	Recent Obs: 2020
<input type="checkbox"/> SO ID: 51160422	Acres: 3,095	Obs Count: 17	Earliest Obs: 2004	Recent Obs: 2018
<input type="checkbox"/> SO ID: 51160858	Acres: 3,095	Obs Count: 6	Earliest Obs: 2014	Recent Obs: 2022
<input type="checkbox"/> SO ID: 51161101	Acres: 3,105	Obs Count: 1	Earliest Obs: 2008	Recent Obs: 2008

Citation for this report:

Montana SOC Occurrences Report

SOC Occurrences for Birds = Bald Eagle

Within Lat/Long: (46.83878,-113.95131) to (46.90232,-114.10672)

Natural Heritage Map Viewer. Montana Natural Heritage Program.

 Retrieved on October 17, 2023, from <https://mtnhp.org/MapView/SOReport.aspx>



MONTANA STATE LIBRARY

NATURAL HERITAGE PROGRAM

mtnhp.org

1201 11th Ave. • P.O. Box 201800 • Helena, MT 59620-1800 • fax 406-444-0266 • phone 406-444-3989



Latitude Longitude
46.84630 -113.99764
46.89484 -114.06045

Summarized by:
013N019W020
(Buffered PLSS Section)



Suggested Citation

Montana Natural Heritage Program. Environmental Summary Report.
for Latitude 46.84630 to 46.89484 and Longitude -113.99764 to -114.06045. Retrieved on 10/17/2023.

The Montana Natural Heritage Program is part of the Montana State Library's Natural Resource Information System. Since 1985, it has served as a neutral and non-regulatory provider of easily accessible information on Montana's species and biological communities to inform all stakeholders in environmental review, permitting, and planning processes. The program is part of the NatureServe network that is composed of over 60 member programs across North America that work to provide current and comprehensive distribution and status information on species and biological communities.

Table of Contents

- [Species Report](#)
- [Structured Surveys](#)
- [Land Cover](#)
- [Wetland and Riparian](#)
- [Land Management](#)
- [Biological Reports](#)
- [Invasive and Pest Species](#)
- [Introduction to Montana Natural Heritage Program](#)
- [Data Use Terms and Conditions](#)
- [Suggested Contacts for Natural Resource Agencies](#)
- [Introduction to Native Species](#)
- [Introduction to Land Cover](#)
- [Introduction to Wetland and Riparian](#)
- [Introduction to Land Management](#)
- [Introduction to Invasive and Pest Species](#)
- [Additional Information Resources](#)

Introduction to Environmental Summary Report

Environmental Summary Reports from the Montana Natural Heritage Program (MTNHP) provide information on species and biological communities to inform all stakeholders in environmental review, permitting, and planning processes. For information on environmental permits in Montana, please see permitting overviews by the [Montana Department of Environmental Quality](#), the [Montana Department of Natural Resources and Conservation](#), the [Index of Environmental Permits for Montana](#) and our [Suggested Contacts for Natural Resource Management Agencies](#). The report for your area of interest consists of introductory and related materials in this PDF and an Excel workbook with worksheets summarizing information managed in the MTNHP databases for: (1) species occurrences; (2) other observed species without species occurrences; (3) other species potentially present based on their range, presence of associated habitats, or predictive distribution model output if available; (4) structured surveys that follow a protocol capable of detecting one or more species; (5) land cover mapped as ecological systems; (6) wetland and riparian mapping; (7) land management categories; and (8) biological reports associated with plant and animal observations. If your area of interest corresponds to a statewide polygon layer (e.g., watersheds, counties, or public land survey sections) information summaries in your report will exactly match those boundaries. However, if your report is for a custom area, users should be aware that summaries do not correspond to the exact boundaries of the polygon they have specified, but instead are a summary across a layer of hexagons intersected by the polygon they specified as shown on the report cover. Summarizing by these hexagons which are one square mile in area and approximately one kilometer in length on each side allows for consistent and rapid delivery of summaries based on a uniform grid that has been used for planning efforts across North America.

In presenting this information, MTNHP is working towards assisting the user with rapidly assessing the known or potential species and biological communities, land management categories, and biological reports associated with the report area. Users are reminded that this information is likely incomplete and may be inaccurate as surveys to document species are lacking in many areas of the state, species' range polygons often include regions of unsuitable habitat, methods of predicting the presence of species or communities are constantly improving, and information is constantly being added and updated in our databases. **Field verification by professional biologists of the absence or presence of species and biological communities in a report area will always be an important obligation of users of our data. Users are encouraged to only use this environmental summary report as a starting point for more in depth analyses and are encouraged to contact state, federal, and tribal resource management agencies for additional data or management guidelines relevant to your efforts. Please see the Appendix for introductory materials to each section of the report, additional information resources, and a list of relevant agency contacts.**

[View in Field Guide](#) [View Predicted Models](#) [View Range Maps](#)

Species of Concern - Native Species Global: G4 State: S2S3 USFWS: C

Delineation Criteria Confirmed breeding area based on the presence of a resident animal of any age/stage. Point observation location is buffered by a minimum distance of 2,000 meters in order to encompass documented travel distances of some butterfly species as well as adjacent habitat likely to support other individuals and otherwise is buffered by the locational uncertainty associated with the observation up to a maximum distance of 10,000 meters. (Last Updated: Sep 22, 2023)

Predicted Models:  79% Optimal (inductive),  21% Moderate (inductive)

B - *Lewis's Woodpecker* (*Melanerpes lewis*) SOC

|22|36|  

[View in Field Guide](#) [View Predicted Models](#) [View Range Maps](#)

Species of Concern - Native Species Global: G4 State: S2B USFWS: MBTA; BCC10; BCC17 USFS: Species of Conservation Concern in Forests (HLC)

BLM: SENSITIVE FWP SWAP: SGCN2 PIF: 2

Delineation Criteria Confirmed breeding area based on the presence of a nest, chicks, or territorial adults during the breeding season. Point observation location is buffered by a minimum distance of 300 meters in order to encompass the likely foraging area used by breeding adults around the nest tree and otherwise is buffered by the locational uncertainty associated with the observation up to a maximum distance of 10,000 meters. (Last Updated: Mar 22, 2023)

Predicted Models:  27% Optimal (inductive),  48% Moderate (inductive),  25% Low (inductive)

B - *Great Blue Heron* (*Ardea herodias*) SOC

|8|172|  

[View in Field Guide](#) [View Predicted Models](#) [View Range Maps](#)

Species of Concern - Native Species Global: G5 State: S3 USFWS: MBTA FWP SWAP: SGCN3

Delineation Criteria Confirmed nesting area buffered by a minimum distance of 6,500 meters in order to be conservative about encompassing the areas commonly used for foraging near the breeding colony and otherwise buffered by the locational uncertainty associated with the observation up to a maximum distance of 10,000 meters. (Last Updated: Jun 27, 2023)

Predicted Models:  13% Optimal (inductive),  30% Moderate (inductive),  57% Low (inductive)

B - *Bald Eagle* (*Haliaeetus leucocephalus*) SSS

|1|180|  

[View in Field Guide](#) [View Predicted Models](#) [View Range Maps](#)

Special Status Species - Native Species Global: G5 State: S4 USFWS: BGEPA; MBTA USFS: Sensitive - Known in Forests (BD, BRT, KOOT, LOLO) BLM: SENSITIVE PIF: 2

Delineation Criteria Confirmed nesting area buffered by a minimum distance of 2,000 meters in order to be conservative about encompassing the breeding territory and area commonly used for renesting. Only nesting observations with a locational uncertainty of 1,000 meters or less will be used to delineate a nesting area. (Last Updated: Sep 05, 2023)

Predicted Models:  3% Optimal (inductive),  46% Moderate (inductive),  48% Low (inductive)

B - *Veery* (*Catharus fuscescens*) SOC

|7|4|  

[View in Field Guide](#) [View Predicted Models](#) [View Range Maps](#)

Species of Concern - Native Species Global: G5 State: S3B USFWS: MBTA BLM: SENSITIVE FWP SWAP: SGCN3 PIF: 2

Delineation Criteria Observations with evidence of breeding activity buffered by a minimum distance of 300 meters in order to be conservative about encompassing home ranges and otherwise buffered by the locational uncertainty associated with the observation up to a maximum distance of 10,000 meters. (Last Updated: Jun 29, 2023)

Predicted Models:  3% Optimal (inductive),  11% Moderate (inductive),  86% Low (inductive)

B - *Evening Grosbeak* (*Coccothraustes vespertinus*) SOC

|120|192|  

[View in Field Guide](#) [View Predicted Models](#) [View Range Maps](#)

Species of Concern - Native Species Global: G5 State: S3 USFWS: MBTA; BCC10 FWP SWAP: SGCN3

Delineation Criteria Confirmed breeding area based on the presence of a nest, chicks, or territorial adults during the breeding season. Point observation location is buffered by a minimum distance of 1,000 meters in order to encompass the maximum foraging distance from nests reported for the species and otherwise is buffered by the locational uncertainty associated with the observation up to a maximum distance of 10,000 meters. (Last Updated: Jun 30, 2023)

Predicted Models:  100% Moderate (inductive)

R - *Western Skink* (*Plestiodon skiltonianus*) SOC

|2|  

[View in Field Guide](#) [View Predicted Models](#) [View Range Maps](#)

Species of Concern - Native Species Global: G5 State: S3 FWP SWAP: SGCN3, SGIN

Delineation Criteria Confirmed breeding area based on the presence of a resident animal of any age. Point observation location is buffered by a minimum distance of 200 meters in order to encompass habitats supporting other individuals in adjacent territories. Otherwise the point observation is buffered by the locational uncertainty associated with the observation up to a maximum distance of 10,000 meters. (Last Updated: Sep 22, 2023)

Predicted Models:  100% Moderate (inductive)

I - *Bombus suckleyi* (Suckley Cuckoo Bumble Bee) SOC

|1|  

[View in Field Guide](#) [View Predicted Models](#) [View Range Maps](#)

Species of Concern - Native Species Global: G2G3 State: S1

Delineation Criteria Confirmed breeding area based on the presence of a resident animal of any age. Point observation location is buffered by a minimum distance of 1700 meters in order to encompass the home range of the individual as well as adjacent habitat likely to support other individuals and otherwise is buffered by the locational uncertainty associated with the observation up to a maximum distance of 10,000 meters. (Last Updated: Jun 22, 2022)

Predicted Models:  100% Moderate (inductive)

M - *Long-legged Myotis* (*Myotis volans*) SOC

|2|  

[View in Field Guide](#) [View Predicted Models](#) [View Range Maps](#)

Species of Concern - Native Species Global: G4G5 State: S3

Delineation Criteria Confirmed area of occupancy based on the documented presence (mistnet captures, definitively identified acoustic recordings, and definitively identified roosting individuals) of adults or juveniles. Point observation location is buffered by a minimum distance of 2,000 meters in order to encompass the average distances traveled from capture locations to roosts in Washington, Oregon, and in the Black Hills of South Dakota and otherwise buffered by the locational uncertainty associated with the observation up to a maximum distance of 10,000 meters. When cave locations are involved, point observations are mapped in the center of a one-square mile hexagon to protect the exact location of the cave entrance as per the Federal Cave Resource Protection Act and associated regulations (U.S. Code Title 16 Chapter 63, Code of Federal Regulations Title 43 Subtitle A Part 37). The outer edges of the hexagon are then buffered by a distance of 2,000 meters and otherwise by the locational uncertainty associated with the observation up to a maximum distance of 10,000 meters. All of the one-square mile hexagons intersecting this buffered area are presented as the Species Occurrence record. (Last Updated: Jul 06, 2023)

Predicted Models:  36% Moderate (inductive),  64% Low (inductive)

B - *Pileated Woodpecker* (*Dryocopus pileatus*) SOC

|43|85|  

[View in Field Guide](#) [View Predicted Models](#) [View Range Maps](#)

Species of Concern - Native Species Global: G5 State: S3 USFWS: MBTA FWP SWAP: SGCN3 PIF: 2

Delineation Criteria Observations with evidence of breeding activity buffered by a minimum distance of 1,500 meters in order to be conservative about encompassing home ranges and otherwise buffered by the locational uncertainty associated with the observation up to a maximum distance of 10,000 meters. (Last Updated: Jun 28, 2023)

Predicted Models:  13% Moderate (inductive)

M - *Hoary Bat* (*Lasius cinereus*) SOC

|2|  

[View in Field Guide](#) [View Predicted Models](#) [View Range Maps](#)

Species of Concern - Native Species Global: G3G4 State: S3B BLM: SENSITIVE FWP SWAP: SGCN3

Delineation Criteria Confirmed area of occupancy based on the documented presence (mistnet captures, definitively identified acoustic recordings, and definitively identified roosting individuals) of adults or juveniles during the active season. Point observation location is buffered by a minimum distance of 3,500 meters in order to be conservative about encompassing the maximum reported foraging distance for the congeneric *Lasiurus borealis* and otherwise buffered by the locational uncertainty associated with the observation up to a maximum distance of 10,000 meters. (Last Updated: Jul 06, 2023)

Predicted Models:  3% Moderate (inductive),  71% Low (inductive)

[View in Field Guide](#) [View Predicted Models](#) [View Range Maps](#)

Species of Concern - Native Species

Global: G5 State: S3

Delineation Criteria Confirmed area of occupancy based on the documented presence (mistnet captures, definitively identified acoustic recordings, and definitively identified roosting individuals) of adults or juveniles. Point observation location is buffered by a minimum distance of 1,000 meters in order to encompass the average distances traveled from capture locations to roosts and between roosts in western Montana, Alberta, and Oregon and otherwise buffered by the locational uncertainty associated with the observation up to a maximum distance of 10,000 meters. When cave locations are involved, point observations are mapped in the center of a one-square mile hexagon to protect the exact location of the cave entrance as per the Federal Cave Resource Protection Act and associated regulations (U.S. Code Title 16 Chapter 63, Code of Federal Regulations Title 43 Subtitle A Part 37). The outer edges of the hexagon are then buffered by a distance of 1,000 meters and otherwise by the locational uncertainty associated with the observation up to a maximum distance of 10,000 meters. All of the one-square mile hexagons intersecting this buffered area are presented as the Species Occurrence record. (Last Updated: Mar 22, 2023)

Predicted Models: 3% Moderate (inductive), 30% Low (inductive)

 V - Isoetes echinospora (*Spiny-spore Quillwort*) SOC

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 90 91 92 93 94 95 96 97 98 99 100

[View in Field Guide](#) [View Predicted Models](#) [View Range Maps](#)

Species of Concern - Native Species

Global: G5 State: S3 Plant Threat Score: **No Known Threats** CCVI: **Less Vulnerable**

Predicted Models: 94% Low (inductive)

 B - Cassin's Finch (*Haemorhous cassinii*) SOC

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 90 91 92 93 94 95 96 97 98 99 100

[View in Field Guide](#) [View Predicted Models](#) [View Range Maps](#)

Species of Concern - Native Species

Global: G5 State: S3 USFWS: **MBTA; BCC10** FWP SWAP: **SGCN3** PIF: 3

Delineation Criteria Observations with evidence of breeding activity buffered by a minimum distance of 300 meters in order to be conservative about encompassing the courtship and foraging distance from nesting areas and otherwise buffered by the locational uncertainty associated with the observation up to a maximum distance of 10,000 meters. (Last Updated: Jun 30, 2023)

Predicted Models: 37% Low (inductive)

 V - Phlox kelseyi var. missoulensis (*Missoula Phlox*) SOC

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 90 91 92 93 94 95 96 97 98 99 100

[View in Field Guide](#) [View Predicted Models](#) [View Range Maps](#)USFWS: **Sensitive - Known in Forests (BD)**

Sensitive - Suspected in Forests (LOLO)

Species of Concern - Native Species

Global: G3 State: S3 **Species of Conservation Concern in Forests (HLC)** CCVI: **Highly Vulnerable**

Delineation Criteria Individual occurrences are generally based upon a discretely mapped area provided by an observer and are not separated by any pre-defined distance. Individual clusters of plants mapped at fine spatial scales (separated by less than approximately 25-50 meters) may be grouped together into one occurrence if they are not separated by distinct areas of habitat or terrain features. Point observations are buffered to encompass any locational uncertainty associated with the observation. (Last Updated: Jun 01, 2023)

Predicted Models: 35% Low (inductive)

 M - Grizzly Bear (*Ursus arctos*) SOC

7 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 90 91 92 93 94 95 96 97 98 99 100

[View in Field Guide](#) [View Range Maps](#)

Species of Concern - Native Species

Global: G4 State: S2 S3 USFWS: **LT** BLM: **THREATENED** FWP SWAP: **SGCN2-3**

Delineation Criteria Species Occurrence polygons represent areas delineated by the U.S. Fish and Wildlife Service (USFWS) that encompass both home ranges and potential transitory movements based on verified sightings. Within these areas, the USFWS wants project proponents to consider whether the species *âœœ* may be presentâœ when evaluating the potential impacts of a project and to work with the USFWS to develop and implement best management practices to minimize or eliminate project effects on the species. (Last Updated: Jul 06, 2023)

 B - American Goshawk (*Accipiter atricapillus*) SOC

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 90 91 92 93 94 95 96 97 98 99 100

[View in Field Guide](#) [View Range Maps](#)

Species of Concern - Native Species

Global: G5 State: S3 USFWS: **MBTA** FWP SWAP: **SGCN3** PIF: 2

Delineation Criteria Confirmed nesting area buffered by a minimum distance of 750 meters in order to encompass the area around the nest known to be defended by adults and otherwise buffered by the locational uncertainty associated with the observation up to a maximum distance of 10,000 meters. (Last Updated: Jan 10, 2023)

 B - Brown Creeper (*Certhia americana*) SOC

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 90 91 92 93 94 95 96 97 98 99 100

[View in Field Guide](#) [View Range Maps](#)

Species of Concern - Native Species

Global: G5 State: S3 USFWS: **MBTA** FWP SWAP: **SGCN3** PIF: 1

Delineation Criteria Observations with evidence of breeding activity buffered by a minimum distance of 300 meters in order to be conservative about encompassing home ranges and otherwise buffered by the locational uncertainty associated with the observation up to a maximum distance of 10,000 meters. (Last Updated: Jun 29, 2023)

 B - Gray-crowned Rosy-Finch (*Leucosticte tephrocotis*) SOC

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 90 91 92 93 94 95 96 97 98 99 100

[View in Field Guide](#) [View Predicted Models](#)

Species of Concern - Native Species

Global: G5 State: S2 USFWS: **MBTA** FWP SWAP: **SGCN2, SGIN**

Delineation Criteria Confirmed breeding area based on the presence of a nest, chicks, or territorial adults during the breeding season. Point observation location is buffered by a minimum distance of 4,000 meters in order to encompass the known foraging distance from nests of the congeneric Black Rosy-Finch and otherwise is buffered by the locational uncertainty associated with the observation up to a maximum distance of 10,000 meters. (Last Updated: Jan 03, 2023)

 B - Great Gray Owl (*Strix nebulosa*) SOC

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 90 91 92 93 94 95 96 97 98 99 100

[View in Field Guide](#) [View Range Maps](#)

Species of Concern - Native Species

Global: G5 State: S3 USFWS: **MBTA** BLM: **SENSITIVE** FWP SWAP: **SGCN3, SGIN** PIF: 3

Delineation Criteria Confirmed nesting area buffered by a minimum distance of 3,200 meters in order to encompass the known foraging distance and area likely to be used for renesting and otherwise buffered by the locational uncertainty associated with the observation up to a maximum distance of 10,000 meters. (Last Updated: Jan 04, 2023)

 B - Pacific Wren (*Troglodytes pacificus*) SOC

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 90 91 92 93 94 95 96 97 98 99 100

[View in Field Guide](#) [View Range Maps](#)

Species of Concern - Native Species

Global: G5 State: S3 USFWS: **MBTA** FWP SWAP: **SGCN3** PIF: 2

Delineation Criteria Observations with evidence of breeding activity buffered by a minimum distance of 300 meters in order to be conservative about encompassing home ranges and otherwise buffered by the locational uncertainty associated with the observation up to a maximum distance of 10,000 meters. (Last Updated: Jun 29, 2023)

 I - *Austrotyla montana* (*A Millipede*) SOC

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 90 91 92 93 94 95 96 97 98 99 100

[View in Field Guide](#) [View Predicted Models](#)

Species of Concern - Native Species

Global: G1 G2 State: S1 S3

Delineation Criteria Confirmed breeding area based on the presence of a resident animal of any age. Point observation location is buffered by a minimum distance of 100 meters in order to encompass the home range of the individual as well as adjacent habitat likely to support other individuals and otherwise is buffered by the locational uncertainty associated with the observation up to a maximum distance of 10,000 meters. (Last Updated: Feb 05, 2008)

 I - *Stygobromus tritus* (*A Subterranean Amphipod*) SOC

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 90 91 92 93 94 95 96 97 98 99 100

[View in Field Guide](#) [View Range Maps](#)

Species of Concern - Native Species

Global: G1 State: S1 S2

Delineation Criteria Confirmed breeding area based on the presence of a resident animal of any age. Point observation location is buffered by a minimum distance of 300 meters in order to encompass the cave system the species is dependent on. (Last Updated: Jan 17, 2008)

[View in Field Guide](#) [View Range Maps](#)**Species of Concern - Native Species**Global: **G5** State: **S3B** USFWS: **MBTA** BLM: **SENSITIVE** FWP SWAP: **SGCN3** PIF: **2****Delineation Criteria** Confirmed breeding area based on the presence of a nest, chicks, or territorial adults during the breeding season. Point observation location is buffered by a minimum distance of 100 meters in order to encompass the maximum territory size reported for the species and otherwise is buffered by the locational uncertainty associated with the observation up to a maximum distance of 10,000 meters. (Last Updated: Jun 29, 2023) **B - Flammulated Owl** (*Psiloscops flammeolus*) SOC

1

1

Not Assessed

S M

[View in Field Guide](#) [View Range Maps](#)**Species of Concern - Native Species**Global: **G4** State: **S3B** USFWS: **MBTA; BCC10** **Species of Conservation Concern in Forests (FLAT, HLC)** BLM: **SENSITIVE**FWP SWAP: **SGCN3** PIF: **1****Delineation Criteria** Confirmed breeding area based on the presence of a nest, chicks, or territorial adults during the breeding season. Point observation location is buffered by a minimum distance of 300 meters in order to encompass the maximum breeding territory size reported for the species and otherwise is buffered by the locational uncertainty associated with the observation up to a maximum distance of 10,000 meters. (Last Updated: Jun 28, 2023) **B - Varied Thrush** (*Ixoreus naevius*) SOC

5

5

Not Assessed

S M

[View in Field Guide](#) [View Range Maps](#)**Species of Concern - Native Species**Global: **G5** State: **S3B** USFWS: **MBTA** FWP SWAP: **SGCN3** PIF: **3****Delineation Criteria** Confirmed breeding area based on the presence of a nest, chicks, or territorial adults during the breeding season. Point observation location is buffered by a minimum distance of 225 meters in order to encompass the reported minimum stand size occupied by breeding pairs and otherwise is buffered by the locational uncertainty associated with the observation up to a maximum distance of 10,000 meters. (Last Updated: Jun 29, 2023) **O - Bat Roost (Non-Cave)** (*Bat Roost (Non-Cave)*) IAH

12

Not Assessed

[View in Field Guide](#)**Important Animal Habitat - Native Species**Global: **GNR** State: **SNR****Delineation Criteria** Confirmed area of occupancy based on the documented presence of adults or juveniles of any bat species at non-cave natural roost sites (e.g. rock outcrops, trees), below ground human created roost sites (e.g. mines), and above ground human created roost sites (e.g., bridges, buildings). Point observation locations are buffered by a distance of 4,500 meters in order to encompass the 95% confidence interval for nightly foraging distance reported for Townsend's Big-eared Bat (a resident Montana bat Species of Concern) and otherwise by the locational uncertainty associated with the observation up to a maximum distance of 10,000 meters. (Last Updated: Oct 22, 2019)

Legend

Model Icons	Habitat Icons	Range Icons	Num Obs
			Count of obs with 'good precision' (<=100m)
			+ indicates additional 'poor precision' obs (1001m-10,000m)


 Latitude 46.84630
 Longitude -113.99764
 46.89484 -114.06045

Native Species

Summarized by: 013N019W020 (Buffered PLSS Section)

Filtered by:

Native Species reports are filtered for Species with MT Status = Species of Concern, Special Status, Important Animal Habitat, Potential SOC

Other Observed Species

		USFWS Sec7	# Obs	Predicted Model	Range
<input type="checkbox"/> B - Hooded Merganser (<i>Lophodytes cucullatus</i>) PSOC	View in Field Guide View Predicted Models View Range Maps	Potential Species of Concern - Native Species	Global: G5 State: S4 USFWS: MBTA FWP SWAP: SGIN PIF: 2		
		Predicted Models:	96% Optimal (inductive), 4% Moderate (inductive)		
<input type="checkbox"/> B - Yellow-billed Cuckoo (<i>Coccyzus americanus</i>) SOC	View in Field Guide View Predicted Models View Range Maps	Species of Concern - Native Species	Global: G5 State: S3B USFWS: PS: LT; MBTA BLM: THREATENED FWP SWAP: SGCN3, SGIN PIF: 2		
		Predicted Models:	80% Optimal (inductive), 20% Moderate (inductive)		
<input type="checkbox"/> B - Rufous Hummingbird (<i>Selasphorus rufus</i>) PSOC	View in Field Guide View Predicted Models View Range Maps	Potential Species of Concern - Native Species	Global: G4 State: S4B USFWS: MBTA; BCC10 PIF: 3		
		Predicted Models:	56% Optimal (inductive), 44% Moderate (inductive)		
<input type="checkbox"/> B - Barrow's Goldeneye (<i>Bucephala islandica</i>) PSOC	View in Field Guide View Predicted Models View Range Maps	Potential Species of Concern - Native Species	Global: G5 State: S4 USFWS: MBTA FWP SWAP: SGIN PIF: 2		
		Predicted Models:	15% Optimal (inductive), 80% Moderate (inductive), 4% Low (inductive)		
<input type="checkbox"/> B - Western Screech-Owl (<i>Megascops kennicottii</i>) PSOC	View in Field Guide View Predicted Models View Range Maps	Potential Species of Concern - Native Species	Global: G4G5 State: S3S4 USFWS: MBTA FWP SWAP: SGIN PIF: 3		
		Predicted Models:	14% Optimal (inductive), 86% Moderate (inductive)		
<input type="checkbox"/> B - American White Pelican (<i>Pelecanus erythrorhynchos</i>) SOC	View in Field Guide View Predicted Models View Range Maps	Species of Concern - Native Species	Global: G4 State: S3B USFWS: MBTA FWP SWAP: SGCN3 PIF: 3		
		Predicted Models:	96% Moderate (inductive), 4% Low (inductive)		
<input type="checkbox"/> B - Black-backed Woodpecker (<i>Picoides arcticus</i>) SOC	View in Field Guide View Predicted Models View Range Maps	Species of Concern - Native Species	Global: G5 State: S3 USFWS: MBTA USFS: Sensitive - Known in Forests (BD, BRT, KOOT, LOLO) BLM: SENSITIVE FWP SWAP: SGCN3 PIF: 1		
		Predicted Models:	26% Moderate (inductive), 71% Low (inductive)		
<input type="checkbox"/> R - Snapping Turtle (<i>Chelydra serpentina</i>) SOC	View in Field Guide View Predicted Models View Range Maps	Species of Concern - Native/Non-native Species - (depends on location or taxa)	Global: G5 State: S3 BLM: SENSITIVE FWP SWAP: SGCN3, SGIN		
		Predicted Models:	26% Moderate (inductive), 60% Low (inductive)		
<input type="checkbox"/> B - Trumpeter Swan (<i>Cygnus buccinator</i>) SOC	View in Field Guide View Predicted Models View Range Maps	Species of Concern - Native Species	Global: G4 State: S3 USFWS: MBTA USFS: Sensitive - Known in Forests (BD) BLM: SENSITIVE FWP SWAP: SGCN3 PIF: 1		
		Predicted Models:	13% Moderate (inductive), 87% Low (inductive)		
<input type="checkbox"/> B - Black-crowned Night-Heron (<i>Nycticorax nycticorax</i>) SOC	View in Field Guide View Predicted Models View Range Maps	Species of Concern - Native Species	Global: G5 State: S3B USFWS: MBTA FWP SWAP: SGCN3 PIF: 3		
		Predicted Models:	13% Moderate (inductive), 28% Low (inductive)		
<input type="checkbox"/> B - Long-billed Curlew (<i>Numenius americanus</i>) SOC	View in Field Guide View Predicted Models View Range Maps	Species of Concern - Native Species	Global: G5 State: S3B USFWS: MBTA; BCC11 BLM: SENSITIVE FWP SWAP: SGCN3 PIF: 2		
		Predicted Models:	6% Moderate (inductive), 94% Low (inductive)		
<input type="checkbox"/> B - Harlequin Duck (<i>Histrionicus histrionicus</i>) SOC	View in Field Guide View Predicted Models View Range Maps	Species of Concern - Native Species	Global: G4 State: S2B USFWS: MBTA USFS: Sensitive - Known in Forests (BD, KOOT, LOLO) FWP SWAP: SGCN2 PIF: 1		
		Predicted Models:	60% Low (inductive)		
<input type="checkbox"/> M - Silver-haired Bat (<i>Lasionycteris noctivagans</i>) PSOC	View in Field Guide View Predicted Models View Range Maps	Potential Species of Concern - Native Species	Global: G3G4 State: S4		
		Predicted Models:	43% Low (inductive)		

<input type="checkbox"/> B - Black-necked Stilt (<i>Himantopus mexicanus</i>) SOC	6	 
View in Field Guide View Predicted Models View Range Maps		
Species of Concern - Native Species Global: G5 State: S3B USFWS: MBTA FWP SWAP: SGCN3 PIF: 3		
Predicted Models:  26% Low (inductive)		
<input type="checkbox"/> B - Horned Grebe (<i>Podiceps auritus</i>) SOC	15	 
View in Field Guide View Predicted Models View Range Maps		
Species of Concern - Native Species Global: G5 State: S3B USFWS: MBTA BLM: SENSITIVE FWP SWAP: SGCN3 PIF: 2		
Predicted Models:  26% Low (inductive)		
<input type="checkbox"/> B - Golden Eagle (<i>Aquila chrysaetos</i>) SOC	9	Not Assessed 
View in Field Guide View Range Maps		
Species of Concern - Native Species Global: G5 State: S3 USFWS: BGEPA; MBTA BLM: SENSITIVE FWP SWAP: SGCN3		
<input type="checkbox"/> I - Rhionaeschna californica (<i>California Darner</i>) PSOC	1	Not Assessed 
View in Field Guide View Range Maps		
Potential Species of Concern - Native Species Global: G5 State: S3S5		
<input type="checkbox"/> B - Black Swift (<i>Cypseloides niger</i>) SOC	2	Not Assessed  
View in Field Guide View Range Maps		
Species of Concern - Native Species Global: G4 State: S1B USFWS: MBTA; BCC10 USFS: Species of Conservation Concern in Forests (FLAT) FWP SWAP: SGCN1, SGIN PIF: 2		
<input type="checkbox"/> B - Caspian Tern (<i>Hydroprogne caspia</i>) SOC	1	Not Assessed 
View in Field Guide View Range Maps		
Species of Concern - Native Species Global: G5 State: S2B USFWS: MBTA BLM: SENSITIVE FWP SWAP: SGCN2 PIF: 2		
<input type="checkbox"/> B - Common Loon (<i>Gavia immer</i>) SOC	4	Not Assessed 
View in Field Guide View Range Maps		
Species of Concern - Native Species Global: G5 State: S3B USFWS: MBTA USFS: Sensitive - Known in Forests (KOOT, LOLO) FWP SWAP: SGCN3 PIF: 1		
<input type="checkbox"/> B - Ferruginous Hawk (<i>Buteo regalis</i>) SOC	12	Not Assessed 
View in Field Guide View Range Maps		
Species of Concern - Native Species Global: G4 State: S3B USFWS: MBTA; BCC17 BLM: SENSITIVE FWP SWAP: SGCN3 PIF: 2		
<input type="checkbox"/> B - Franklin's Gull (<i>Leucophaeus pipixcan</i>) SOC	3	Not Assessed 
View in Field Guide View Range Maps		
Species of Concern - Native Species Global: G5 State: S3B USFWS: MBTA; BCC10; BCC11; BCC17 BLM: SENSITIVE FWP SWAP: SGCN3 PIF: 2		
<input type="checkbox"/> B - Tennessee Warbler (<i>Leiothlypis peregrina</i>) PSOC	1	Not Assessed 
View in Field Guide View Range Maps		
Potential Species of Concern - Native Species Global: G5 State: S3S4B USFWS: MBTA		
<input type="checkbox"/> B - White-faced Ibis (<i>Plegadis chihi</i>) SOC	2	Not Assessed 
View in Field Guide View Range Maps		
Species of Concern - Native Species Global: G5 State: S3B USFWS: MBTA BLM: SENSITIVE FWP SWAP: SGCN3 PIF: 2		
<input type="checkbox"/> B - Burrowing Owl (<i>Athene cunicularia</i>) SOC	3	Not Assessed
View in Field Guide		
Species of Concern - Native Species Global: G4 State: S3B USFWS: MBTA; BCC17 BLM: SENSITIVE FWP SWAP: SGCN3 PIF: 1		
<input type="checkbox"/> B - Sagebrush Sparrow (<i>Artemiospiza nevadensis</i>) SOC	2	Not Assessed
View in Field Guide		
Species of Concern - Native Species Global: G5 State: S3B USFWS: MBTA BLM: SENSITIVE FWP SWAP: SGCN3		

Legend

Model Icons	Habitat Icons	Range Icons	Num Obs
Suitable (native range)	Common	Native / Year-round	Count of obs with 'good precision' (<=100m)
Optimal Suitability	Occasional	Summer	+ indicates additional 'poor precision' obs (1001m-10,000m)
Moderate Suitability		Winter	
Low Suitability		Migratory	
Suitable (introduced range)		Non-native	
		Historical	


Latitude 46.84630
Longitude -113.99764
46.89484 -114.06045

Native Species














Summarized by: **013N019W020 (Buffered PLSS Section)**

Filtered by:

Native Species reports are filtered for Species with MT Status = Species of Concern, Special Status, Important Animal Habitat, Potential SOC

Other Potential Species

		USFWS Sec7	Predicted Model	Range
<input type="checkbox"/> V - <i>Camissonia andina</i> (<i>Obscure Evening-primrose</i>) SOC	View in Field Guide View Predicted Models View Range Maps			Y
Species of Concern - Native Species Global: G4 State: S2 Plant Threat Score: No Known Threats				
Predicted Models: 5% Suitable (native range) (deductive)				
<input type="checkbox"/> V - <i>Rotala ramosior</i> (<i>Toothcup</i>) SOC	View in Field Guide View Predicted Models View Range Maps			Y
Species of Concern - Native Species Global: G5 State: S1S2 Plant Threat Score: No Known Threats CCVI: Highly Vulnerable				
Predicted Models: 1% Suitable (native range) (deductive)				
<input type="checkbox"/> M - <i>Western Spotted Skunk</i> (<i>Spilogale gracilis</i>) PSOC	View in Field Guide View Predicted Models View Range Maps		Y	Y
Potential Species of Concern - Native Species Global: G5 State: SU FWP SWAP: SGIN				
Predicted Models: 91% Optimal (inductive), 9% Moderate (inductive)				
<input type="checkbox"/> B - <i>Broad-tailed Hummingbird</i> (<i>Selasphorus platycercus</i>) PSOC	View in Field Guide View Predicted Models View Range Maps			S M
Potential Species of Concern - Native Species Global: G5 State: S4B USFWS: MBTA; BCC10 FWP SWAP: SGIN				
Predicted Models: 91% Optimal (inductive), 9% Moderate (inductive)				
<input type="checkbox"/> V - <i>Dichanthelium acuminatum</i> (<i>Panic Grass</i>) SOC	View in Field Guide View Predicted Models View Range Maps		Y	Y
Species of Concern - Native Species Global: G5 State: S2S3 Plant Threat Score: Unknown				
Predicted Models: 67% Optimal (inductive), 33% Moderate (inductive)				
<input type="checkbox"/> V - <i>Impatiens aurella</i> (<i>Pale-yellow Jewel-weed</i>) SOC	View in Field Guide View Predicted Models View Range Maps		Y	Y
Species of Concern - Native Species Global: G4 State: S3 Plant Threat Score: No Known Threats				
Predicted Models: 59% Optimal (inductive), 41% Moderate (inductive)				
<input type="checkbox"/> V - <i>Carex scoparia</i> (<i>Pointed Broom Sedge</i>) SOC	View in Field Guide View Predicted Models View Range Maps		Y	Y
Species of Concern - Native Species Global: G5 State: S1S2 Plant Threat Score: No Known Threats				
Predicted Models: 54% Optimal (inductive), 46% Moderate (inductive)				
<input type="checkbox"/> V - <i>Wolffia columbiana</i> (<i>Columbia Water-meal</i>) SOC	View in Field Guide View Predicted Models View Range Maps		Y	Y
Species of Concern - Native Species Global: G5 State: S2S3 Plant Threat Score: No Known Threats				
Predicted Models: 42% Optimal (inductive), 58% Moderate (inductive)				
<input type="checkbox"/> I - <i>Zumatrichia notosa</i> (<i>A Caddisfly</i>) SOC	View in Field Guide View Predicted Models View Range Maps		Y	Y
Species of Concern - Native Species Global: G2G4 State: S3				
Predicted Models: 11% Optimal (inductive), 26% Moderate (inductive), 24% Low (inductive)				
<input type="checkbox"/> V - <i>Cypripedium parviflorum</i> (<i>Small Yellow Lady's-slipper</i>) PSOC	View in Field Guide View Predicted Models View Range Maps		Y	Y
Potential Species of Concern - Native Species Global: G5 State: S3S4 USFS: Sensitive - Known in Forests (KOOT, LOLO) Sensitive - Suspected in Forests (BRT) Species of Conservation Concern in Forests (CG, HLC)				
Predicted Models: 1% Optimal (inductive), 35% Moderate (inductive), 64% Low (inductive)				
<input type="checkbox"/> V - <i>Utricularia intermedia</i> (<i>Flatleaf Bladderwort</i>) SOC	View in Field Guide View Predicted Models View Range Maps		Y	Y
Species of Concern - Native Species Global: G5 State: S2 USFS: Sensitive - Known in Forests (KOOT) Plant Threat Score: No Known Threats				
Predicted Models: 96% Moderate (inductive), 4% Low (inductive)				
<input type="checkbox"/> V - <i>Juncus covillei</i> (<i>Coville's Rush</i>) SOC	View in Field Guide View Predicted Models View Range Maps		Y	Y
Species of Concern - Native Species Global: G5 State: S2S3 Plant Threat Score: No Known Threats				
Predicted Models: 72% Moderate (inductive), 28% Low (inductive)				
<input type="checkbox"/> V - <i>Botrychium lineare</i> (<i>Linearleaf Moonwort</i>) SOC	View in Field Guide View Predicted Models View Range Maps		Y	Y
Species of Concern - Native Species Global: G3 State: S1S2 CCVI: Less Vulnerable				
Predicted Models: 64% Moderate (inductive), 5% Low (inductive)				

<input type="checkbox"/> M - North American Porcupine (<i>Erethizon dorsatum</i>) PSOC	  
View in Field Guide View Predicted Models View Range Maps	
Potential Species of Concern - Native Species	Global: G5 State: S3S4 FWP SWAP: SGIN
Predicted Models:	 51% Moderate (inductive),  49% Low (inductive)
<input type="checkbox"/> V - Eleocharis rostellata (Beaked Spikerush) SOC	  
View in Field Guide View Predicted Models View Range Maps	
Species of Concern - Native Species	Global: G5 State: S3 USFS: Species of Conservation Concern in Forests (CG, FLAT, HLC) Plant Threat Score: Unknown
CCVI: Less Vulnerable	
Predicted Models:	 33% Moderate (inductive),  13% Low (inductive)
<input type="checkbox"/> V - Ligusticum verticillatum (Idaho Lovage) SOC	  
View in Field Guide View Predicted Models View Range Maps	
Species of Concern - Native Species	Global: G4G5 State: S3 Plant Threat Score: No Known Threats
Predicted Models:	 25% Moderate (inductive),  23% Low (inductive)
<input type="checkbox"/> V - Draba densifolia (Dense-leaf Draba) SOC	  
View in Field Guide View Predicted Models View Range Maps	
Species of Concern - Native Species	Global: G5 State: S2 USFS: Species of Conservation Concern in Forests (CG, HLC) Plant Threat Score: Low
CCVI: Moderately Vulnerable	
Predicted Models:	 20% Moderate (inductive),  39% Low (inductive)
<input type="checkbox"/> B - Meesia triquetra (Meesia Moss) SOC	  
View in Field Guide View Predicted Models View Range Maps	
Species of Concern - Native Species	USFS: Sensitive - Known in Forests (BRT, KOOT) Sensitive - Suspected in Forests (LOLO)
Global: G5 State: S2 Species of Conservation Concern in Forests (CG, FLAT)	
Predicted Models:	 13% Moderate (inductive),  82% Low (inductive)
<input type="checkbox"/> V - Carex cracae (Crake's Sedge) SOC	  
View in Field Guide View Predicted Models View Range Maps	
Species of Concern - Native Species	Global: G5 State: S2S3 Plant Threat Score: Low
Predicted Models:	 13% Moderate (inductive),  41% Low (inductive)
<input type="checkbox"/> A - Western Toad (<i>Anaxyrus boreas</i>) SOC	  
View in Field Guide View Predicted Models View Range Maps	
Species of Concern - Native Species	Global: G4 State: S2 USFS: Sensitive - Known in Forests (BD, BRT, KOOT, LOLO) BLM: SENSITIVE FWP SWAP: SGCN2
Predicted Models:	 13% Moderate (inductive),  22% Low (inductive)
<input type="checkbox"/> V - Centunculus minimus (Chaffweed) SOC	  
View in Field Guide View Predicted Models View Range Maps	
Species of Concern - Native Species	Global: G5 State: S2 Plant Threat Score: No Known Threats
Predicted Models:	 13% Moderate (inductive),  21% Low (inductive)
<input type="checkbox"/> I - Oreohelix haydeni (Lyrate Mountainsnail) SOC	  
View in Field Guide View Predicted Models View Range Maps	
Species of Concern - Native Species	Global: G2 State: S1S3
Predicted Models:	 11% Moderate (inductive),  89% Low (inductive)
<input type="checkbox"/> B - Ovenbird (<i>Seiurus aurocapilla</i>) PSOC	    
View in Field Guide View Predicted Models View Range Maps	
Potential Species of Concern - Native Species	Global: G5 State: S4B USFWS: MBTA PIF: 3
Predicted Models:	 3% Moderate (inductive),  97% Low (inductive)
<input type="checkbox"/> V - Elodea bifoliata (Long-sheath Waterweed) SOC	  
View in Field Guide View Predicted Models View Range Maps	
Species of Concern - Native Species	Global: G4G5 State: S2? Plant Threat Score: No Known Threats
Predicted Models:	 3% Moderate (inductive),  93% Low (inductive)
<input type="checkbox"/> V - Psilocarphus brevissimus (Dwarf woolly-heads) SOC	  
View in Field Guide View Predicted Models View Range Maps	
Species of Concern - Native Species	Global: G4 State: S2S3 USFS: Sensitive - Known in Forests (KOOT) Plant Threat Score: No Known Threats
Predicted Models:	 3% Moderate (inductive),  72% Low (inductive)
<input type="checkbox"/> M - Little Brown Myotis (<i>Myotis lucifugus</i>) SOC	  
View in Field Guide View Predicted Models View Range Maps	
Species of Concern - Native Species	Global: G3G4 State: S3 FWP SWAP: SGCN3
Predicted Models:	 3% Moderate (inductive),  52% Low (inductive)
<input type="checkbox"/> V - Stipa lettermanii (Letterman's Needlegrass) SOC	  
View in Field Guide View Predicted Models View Range Maps	
Species of Concern - Native Species	Global: G5 State: S1S3 USFS: Species of Conservation Concern in Forests (HLC) Plant Threat Score: No Known Threats
Predicted Models:	 1% Moderate (inductive),  11% Low (inductive)
<input type="checkbox"/> V - Drosera rotundifolia (Roundleaf Sundew) PSOC	  
View in Field Guide View Predicted Models View Range Maps	
Potential Species of Concern - Native Species	Global: G5 State: S3S4 Plant Threat Score: Unknown
Predicted Models:	 1% Moderate (inductive),  3% Low (inductive)
<input type="checkbox"/> R - Northern Alligator Lizard (<i>Elgaria coerulea</i>) SOC	  
View in Field Guide View Predicted Models View Range Maps	
Species of Concern - Native Species	Global: G5 State: S3 FWP SWAP: SGCN3, SGIN
Predicted Models:	 100% Low (inductive)

<input type="checkbox"/> V - <i>Stellaria crassifolia</i> (<i>Fleshy Stitchwort</i>) SOC	View in Field Guide	View Predicted Models	View Range Maps									
Species of Concern - Native Species	Global: G5	State: S2	Plant Threat Score: No Known Threats									
Predicted Models:	 100% Low (inductive)											
<input type="checkbox"/> B - <i>Common Poorwill</i> (<i>Phalaenoptilus nuttallii</i>) PSOC	View in Field Guide	View Predicted Models	View Range Maps									
Potential Species of Concern - Native Species	Global: G5	State: S4B	USFWS: MBTA	FWP SWAP: SGIN	PIF: 3							
Predicted Models:	 100% Low (inductive)											
<input type="checkbox"/> V - <i>Ranunculus hyperboreus</i> (<i>High Northern Buttercup</i>) PSOC	View in Field Guide	View Predicted Models	View Range Maps									
Potential Species of Concern - Native Species	Global: G5	State: S3S4										
Predicted Models:	 74% Low (inductive)											
<input type="checkbox"/> M - <i>Canada Lynx</i> (<i><i>Lynx canadensis</i></i>) SOC	View in Field Guide	View Predicted Models	View Range Maps									
Species of Concern - Native Species	Global: G5	State: S3	USFWS: LT; CH	BLM: THREATENED	FWP SWAP: SGCN3							
Predicted Models:	 69% Low (inductive)											
<input type="checkbox"/> V - <i>Erigeron linearis</i> (<i>Linear-leaf Fleabane</i>) SOC	View in Field Guide	View Predicted Models	View Range Maps									
Species of Concern - Native Species	Global: G5	State: S2	Plant Threat Score: Low	CCVI: Less Vulnerable								
Predicted Models:	 31% Low (inductive)											
<input type="checkbox"/> M - <i>Hoary Marmot</i> (<i><i>Marmota caligata</i></i>) PSOC	View in Field Guide	View Predicted Models	View Range Maps									
Potential Species of Concern - Native Species	Global: G5	State: S3S4	FWP SWAP: SGIN									
Predicted Models:	 29% Low (inductive)											
<input type="checkbox"/> B - <i>Black Tern</i> (<i><i>Chlidonias niger</i></i>) SOC	View in Field Guide	View Predicted Models	View Range Maps									
Species of Concern - Native Species	Global: G4G5	State: S3B	USFWS: MBTA; BCC10; BCC11; BCC17	BLM: SENSITIVE	FWP SWAP: SGCN3	PIF: 2						
Predicted Models:	 26% Low (inductive)											
<input type="checkbox"/> F - <i>Lake Trout</i> (<i><i>Salvelinus namaycush</i></i>) SOC	View in Field Guide	View Predicted Models	View Range Maps									
Species of Concern - Native/Non-native Species - (depends on location or taxa)	Global: G5	State: S2	FWP SWAP: SGCN2									
Predicted Models:	 55% Suitable (introduced range) (deductive)											
<input type="checkbox"/> M - <i>Wolverine</i> (<i><i>Gulo gulo</i></i>) SOC	View in Field Guide	View Range Maps										
Species of Concern - Native Species	Global: G4	State: S3	USFS: Sensitive - Known in Forests (BD, BRT, KOOT, LOLO)	BLM: SENSITIVE	FWP SWAP: SGCN3							
						 Not Assessed						

Structured Surveys

Summarized by: **013N019W020 (Buffered PLSS Section)**

The Montana Natural Heritage Program (MTNHP) records information on the locations where more than 80 different types of well-defined repeatable survey protocols capable of detecting an animal species or suite of animal species have been conducted by state, federal, tribal, university, or private consulting biologists. Examples of structured survey protocols tracked by MTNHP include: visual encounter and dip net surveys for pond breeding amphibians, point counts for birds, call playback surveys for selected bird species, visual surveys of migrating raptors, kick net stream reach surveys for macroinvertebrates, visual encounter cover object surveys for terrestrial mollusks, bat acoustic or mist net surveys, pitfall and/or snap trap surveys for small terrestrial mammals, track or camera trap surveys for large mammals, and trap surveys for turtles. Whenever possible, photographs of survey locations are stored in MTNHP databases.



MTNHP does not typically manage information on structured surveys for plants; surveys for invasive species may be a future exception.

Within the report area you have requested, structured surveys are summarized by the number of each type of structured survey protocol that has been conducted, the number of species detections/observations resulting from these surveys, and the most recent year a survey has been conducted.

B-Colonial-nesting Waterbirds <i>(Colonial-nesting Waterbird Surveys)</i>	Survey Count: 2	Obs Count: 2	Recent Survey: 2011
B-Great Blue Heron Rookery <i>(Great Blue Heron Rookery)</i>	Survey Count: 1	Obs Count: 1	Recent Survey: 2021
B-Point Count <i>(Bird Point Count)</i>	Survey Count: 42	Obs Count: 339	Recent Survey: 1994
B-Raptor nest <i>(Raptor Nest Survey)</i>	Survey Count: 2	Obs Count: 2	Recent Survey: 2009
E-Eastern Heath Snail <i>(Eastern Heath Snail Survey)</i>	Survey Count: 3	Obs Count: 3	Recent Survey: 2012
E-Eurasian Water-milfoil Rake <i>(Rake tows/pulls for Eurasian Water-milfoil)</i>	Survey Count: 6	Obs Count: 6	Recent Survey: 2012
E-Kicknet <i>(Kicknet Collection Survey for Invasive Mussels and Snails)</i>	Survey Count: 3	Obs Count: 1	Recent Survey: 2022
E-Noxious Weed, Road-based <i>(Noxious Weed Road-based Visual Surveys)</i>	Survey Count: 23	Obs Count: 14	Recent Survey: 2003
E-Visual Aquatic Invasives <i>(Visual Encounter Surveys for Aquatic Invasives on Shorelines or Underwater)</i>	Survey Count: 5	Obs Count: 5	Recent Survey: 2022
F-Fish Other Survey <i>(Fish Other Survey (FWP Survey Type))</i>	Survey Count: 1	Obs Count: 1	Recent Survey: 1951
I-Odonates/Butterfly VES <i>(Visual Encounter Survey for Damselfly/Dragonfly/Butterfly)</i>	Survey Count: 2	Obs Count: 2	Recent Survey: 1956
M-Bat Roost (Active Season) <i>(Bat Roost (Active Season) Survey)</i>	Survey Count: 2	Obs Count: 1	Recent Survey: 2014
P-FIA Ecosystem <i>(USFS FIA Ecosystem Survey)</i>	Survey Count: 1	Obs Count: 5	Recent Survey: 2003

Land Cover

Summarized by: **013N019W020 (Buffered PLSS Section)**



Human Land Use Developed

Other Roads


29% (1,644 Acres) County, city and or rural roads generally open to motor vehicles.



Human Land Use Developed

Low Intensity Residential

25% (1,442 Acres) Includes areas with a mixture of constructed materials and vegetation. Impervious surfaces account for 20-50% of total cover. These areas most commonly include single-family housing units in rural and suburban areas. Paved roadways may be classified into this category.



Human Land Use Developed

Commercial / Industrial

16% (899 Acres) Businesses, industrial parks, hospitals, airports; utilities in commercial/industrial areas.



Human Land Use Developed

Developed, Open Space

7% (379 Acres) Vegetation (primarily grasses) planted in developed settings for recreation, erosion control, or aesthetic purposes. Impervious surfaces account for less than 20% of total cover. This category often includes highway and railway rights of way and graveled rural roads.



Human Land Use Developed

High Intensity Residential

6% (333 Acres) Includes areas with a mixture of constructed materials and vegetation. Impervious surfaces account for 50-80% of the total cover. These areas most commonly include single-family housing units in urban areas. Paved roadways, parking lots, and other large impervious surfaces may be classified into this category.



Wetland and Riparian Systems

Floodplain and Riparian

4% (223 Acres)

Northern Rocky Mountain Lower Montane Riparian Woodland and Shrubland

This ecological system is found throughout the Rocky Mountain and Colorado Plateau regions. In Montana, sites occur at elevations of 609-1,219 meters (2,000-4,000 feet) west of the Continental Divide. East of the Continental Divide, this system ranges up to 1,676 meters (5,500 feet). It generally comprises a mosaic of multiple communities that are tree-dominated with a diverse shrub component. It is dependent on a natural hydrologic regime with annual to episodic flooding, so it is usually found within the flood zone of rivers, on islands, sand or cobble bars, and along streambanks. It can form large, wide occurrences on mid-channel islands in larger rivers, or narrow bands on small, rocky canyon tributaries and well-drained benches. It is also typically found in backwater channels and other perennially wet but less scoured sites, such as floodplains, swales and irrigation ditches. In some locations, occurrences extend into moderately high intermountain basins where the adjacent vegetation is sage steppe. Black cottonwood (*Populus balsamifera* ssp. *trichocarpa*) is the key indicator species. Other dominant trees may include boxelder maple (*Acer negundo*), narrowleaf cottonwood (*Populus angustifolia*), eastern cottonwood (*Populus deltoides*), Douglas-fir (*Pseudotsuga menziesii*), peachleaf willow (*Salix amygdaloides*), or Rocky Mountain juniper (*Juniperus scopulorum*). Dominant shrubs include Rocky Mountain maple (*Acer glabrum*), thinleaf alder (*Alnus incana*), river birch (*Betula occidentalis*), redosier dogwood (*Cornus sericea*), hawthorne (*Crataegus* species), chokecherry (*Prunus virginiana*), skunkbush sumac (*Rhus trilobata*), willows (*Salix* species), rose (*Rosa* species), silver buffaloberry (*Shepherdia argentea*), or snowberry (*Symphoricarpos* species).



Wetland and Riparian Systems

Open Water

4% (215 Acres)

Open Water

All areas of open water, generally with less than 25% cover of vegetation or soil



Human Land Use

Agriculture

3% (200 Acres)

Cultivated Crops

These areas used for the production of crops, such as corn, soybeans, small grains, sunflowers, vegetables, and cotton, typically on an annual cycle. Agricultural plant cover is variable depending on season and type of farming. Other areas include more stable land cover of orchards and vineyards.



Grassland Systems

Montane Grassland

3% (196 Acres)

Rocky Mountain Lower Montane, Foothill, and Valley Grassland

This grassland system of the northern Rocky Mountains is found at lower montane to foothill elevations in mountains and valleys throughout Montana. These grasslands are floristically similar to Big Sagebrush Steppe but are defined by shorter summers, colder winters, and young soils derived from recent glacial and alluvial material. They are found at elevations from 548 - 1,650 meters (1,800-5,413 feet). In the lower montane zone, they range from small meadows to large open parks surrounded by conifers; below the lower treeline, they occur as extensive foothill and valley grasslands. Soils are relatively deep, fine-textured, often with coarse fragments, and non-saline. Microphytic crust may be present in high-quality occurrences. This system is typified by cool-season perennial bunch grasses and forbs (>25% cover, with a sparse shrub cover (<10%). Rough fescue (*Festuca campestris*) is dominant in the northwestern portion of the state and Idaho fescue (*Festuca idahoensis*) is dominant or co-dominant throughout the range of the system. Bluebunch wheatgrass (*Pseudoroegneria spicata*) occurs as a co-dominant throughout the range as well, especially on xeric sites. Western wheatgrass (*Pascopyrum smithii*) is consistently present, often with appreciable coverage (>10%) in lower elevation occurrences in western Montana and virtually always present, with relatively high coverages (>25%), on the edge of the Northwestern Great Plains region. Species diversity ranges from a high of more than 50 per 400 square meter plot on mesic sites to 15 (or fewer) on xeric and disturbed sites. Most occurrences have at least 25 vascular species present. Farmland conversion, noxious species invasion, fire suppression, heavy grazing and oil and gas development are major threats to this system.

Additional Limited Land Cover

1% (69 Acres) [Railroad](#)

1% (64 Acres) [Major Roads](#)

1% (61 Acres) [Alpine-Montane Wet Meadow](#)

<1% (28 Acres) [Interstate](#)

<1% (9 Acres) [Introduced Upland Vegetation - Annual and Biennial Forland](#)


<1% (2 Acres) [Rocky Mountain Dry-Mesic Montane Mixed Conifer Forest](#)

<1% (1 Acres) [Rocky Mountain Ponderosa Pine Woodland and Savanna](#)

<1% (0 Acres) [Emergent Marsh](#)



Wetland and Riparian

 Summarized by: **013N019W020 (Buffered PLSS Section)**


Wetland and Riparian Mapping

[Explain !\[\]\(0b0636dbae614f97346d733ac650473d_img.jpg\)](#)

P - Palustrine

█ UB - Unconsolidated Bottom	
F - Semipermanently Flooded	<1 Acres
x - Excavated	<1 Acres PUBFx

P - Palustrine, UB - Unconsolidated Bottom
 Wetlands where mud, silt or similar fine particles cover at least 25% of the bottom, and where vegetation cover is less than 30%.

█ AB - Aquatic Bed	
F - Semipermanently Flooded	59 Acres

P - Palustrine, AB - Aquatic Bed
 Wetlands with vegetation growing on or below the water surface for most of the growing season.

(no modifier)	2 Acres PABF
h - Diked/Impounded	3 Acres PABFh
x - Excavated	54 Acres PABFx

G - Intermittently Exposed	5 Acres
x - Excavated	5 Acres PABGx

P - Palustrine, AB - Aquatic Bed
 Wetlands with vegetation growing on or below the water surface for most of the growing season.

K - Artificially Flooded	1 Acres
x - Excavated	1 Acres PABKx

█ US - Unconsolidated Shore	
A - Temporarily Flooded	<1 Acres

P - Palustrine, US - Unconsolidated Shore
 Wetlands with less than 75% areal cover of stones, boulders, or bedrock. AND with less than 30% vegetative cover AND the wetland is irregularly exposed due to seasonal or irregular flooding and subsequent drying.

x - Excavated	<1 Acres PUSAx
---------------	--------------------------

█ EM - Emergent	
A - Temporarily Flooded	16 Acres

P - Palustrine, EM - Emergent
 Wetlands with erect, rooted herbaceous vegetation present during most of the growing season.

(no modifier)	16 Acres PEMA
C - Seasonally Flooded	4 Acres

(no modifier)	4 Acres PEMC
F - Semipermanently Flooded	1 Acres

(no modifier)	1 Acres PEMF
---------------	---------------------

█ SS - Scrub-Shrub	
A - Temporarily Flooded	18 Acres

P - Palustrine, SS - Scrub-Shrub
 Wetlands dominated by woody vegetation less than 6 meters (20 feet) tall. Woody vegetation includes tree saplings and trees that are stunted due to environmental conditions.

(no modifier)	18 Acres PSSA
C - Seasonally Flooded	<1 Acres

R - Riverine (Rivers)**2 - Lower Perennial**

█ UB - Unconsolidated Bottom	R - Riverine (Rivers), 2 - Lower Perennial, UB - Unconsolidated Bottom Stream channels where the substrate is at least 25% mud, silt or other fine particles.		
H - Permanently Flooded (no modifier)	45 Acres	45 Acres	R2UBH
█ US - Unconsolidated Shore	R - Riverine (Rivers), 2 - Lower Perennial, US - Unconsolidated Shore Shorelines with less than 75% areal cover of stones, boulders, or bedrock and less than 30% vegetation cover. The area is also irregularly exposed due to seasonal or irregular flooding and subsequent drying.		
A - Temporarily Flooded (no modifier)	3 Acres	3 Acres	R2USA

3 - Upper Perennial

█ UB - Unconsolidated Bottom	R - Riverine (Rivers), 3 - Upper Perennial, UB - Unconsolidated Bottom Stream channels where the substrate is at least 25% mud, silt or other fine particles.		
G - Intermittently Exposed (no modifier)	81 Acres	81 Acres	R3UBG
█ US - Unconsolidated Shore	R - Riverine (Rivers), 3 - Upper Perennial, US - Unconsolidated Shore Shorelines with less than 75% areal cover of stones, boulders, or bedrock and less than 30% vegetation cover. The area is also irregularly exposed due to seasonal or irregular flooding and subsequent drying.		
A - Temporarily Flooded (no modifier)	17 Acres	17 Acres	R3USA
C - Seasonally Flooded (no modifier)	1 Acres	1 Acres	R3USC

4 - Intermittent

█ SB - Stream Bed	R - Riverine (Rivers), 4 - Intermittent, SB - Stream Bed Active channel that contains periodic water flow.		
C - Seasonally Flooded x - Excavated	11 Acres	11 Acres	R4SBCx

Rp - Riparian**1 - Lotic**


█ SS - Scrub-Shrub (no modifier)	24 Acres	Rp1SS	Rp - Riparian, 1 - Lotic, SS - Scrub-Shrub This type of riparian area is dominated by woody vegetation that is less than 6 meters (20 feet) tall. Woody vegetation includes tree saplings and trees that are stunted due to environmental conditions.
█ FO - Forested (no modifier)	132 Acres	Rp1FO	Rp - Riparian, 1 - Lotic, FO - Forested This riparian class has woody vegetation that is greater than 6 meters (20 feet) tall.
█ EM - Emergent (no modifier)	25 Acres	Rp1EM	Rp - Riparian, 1 - Lotic, EM - Emergent Riparian areas that have erect, rooted herbaceous vegetation during most of the growing season.

2 - Lentic

█ SS - Scrub-Shrub (no modifier)	1 Acres	Rp2SS	Rp - Riparian, 2 - Lentic, SS - Scrub-Shrub This type of riparian area is dominated by woody vegetation that is less than 6 meters (20 feet) tall. Woody vegetation includes tree saplings and trees that are stunted due to environmental conditions.
█ FO - Forested (no modifier)	<1 Acres	Rp2FO	Rp - Riparian, 2 - Lentic, FO - Forested This riparian class has woody vegetation that is greater than 6 meters (20 feet) tall.


 Latitude 46.84630
 Longitude -113.99764
 46.89484
 -114.06045

Land Management

 Summarized by: **013N019W020 (Buffered PLSS Section)**


Land Management Summary

[Explain](#)

	Ownership	Tribal	Easements	Other Boundaries (possible overlap)
Public Lands	424 Acres (7%)			
Federal	19 Acres (<1%)			
US Forest Service	14 Acres (<1%)			
USFS Owned	14 Acres (<1%)			
US Government	5 Acres (<1%)			
US Government Owned	5 Acres (<1%)			
State	44 Acres (1%)			
Montana Department of Transportation	44 Acres (1%)			
MTDOT Owned	44 Acres (1%)			
Local	361 Acres (6%)			
Local Government	361 Acres (6%)			
Local Government Owned	361 Acres (6%)			
Conservation Easements	2 Acres (<1%)			
State & Local	2 Acres (<1%)			
City Government	2 Acres (<1%)			
Private Lands or Unknown Ownership	5,338 Acres (93%)			

Biological Reports

Summarized by: **013N019W020 (Buffered PLSS Section)**

Within the report area you have requested, citations for all reports and publications associated with plant or animal observations in Montana Natural Heritage Program (MTNHP) databases are listed and, where possible, links to the documents are included.

The MTNHP plans to include reports associated with terrestrial and aquatic communities in the future as allowed for by staff resources. If you know of reports or publications associated with species or biological communities within the report area that are not shown in this report, please let us know: mtnhp@mt.gov

- Burleigh, T.D. 1951. ***Spring migration***. Audubon Field Notes 5:266-268.
- Hand, R.L. 1969. ***A distributional checklist of the birds of western Montana***. Unpublished. Available at Mansfield Library, University of Montana, Missoula.
- Hansen, R.M. 1962. ***Movements and survival of Thomomys talpoides in a mima-mound habitat***. Ecology 43(1):151-154.
- Kinsella, J. M. 1966. ***Helminths of Microtinae in western Montana***. M.A. thesis. University of Montana, Missoula. 61 pp.
- Missoula County Weed District. ***Geodatabases with sample site location data related to AIS surveys beginning in 2011 on waterbodies in western Montana***

Legend

Model Icons	Habitat Icons	Range Icons	Num Obs
			Count of obs with 'good precision' (<=1000m)
			+ indicates additional 'poor precision' obs (1001m-10,000m)


Latitude 46.84630
Longitude -113.99764
46.89484 -114.06045

Invasive and Pest Species

Summarized by: 013N019W020 (Buffered PLSS Section)

Aquatic Invasive Species
 V - *Iris pseudacorus* (Yellowflag Iris) N2A/AIS

[View in Field Guide](#) [View Predicted Models](#) [View Range Maps](#)

Noxious Weed: Priority 2A - Aquatic Invasive Species - Non-native Species Global: GNR State: SNA

Predicted Models: 100% Optimal (inductive)

# Obs	Predicted Model	Range
29		

 A - American Bullfrog (*Lithobates catesbeianus*) AIS

[View in Field Guide](#) [View Predicted Models](#) [View Range Maps](#)

Aquatic Invasive Species - Non-native Species Global: G5 State: SNA

Predicted Models: 26% Optimal (inductive), 43% Moderate (inductive), 31% Low (inductive)

--	--	--

 V - *Butomus umbellatus* (Flowering-rush) N2A/AIS

[View in Field Guide](#) [View Predicted Models](#) [View Range Maps](#)

Noxious Weed: Priority 2A - Aquatic Invasive Species - Non-native Species Global: G5 State: SNA

Predicted Models: 11% Optimal (inductive), 23% Moderate (inductive), 26% Low (inductive)

--	--	--

 V - *Potamogeton crispus* (Curly-leaf Pondweed) N2B/AIS

[View in Field Guide](#) [View Predicted Models](#) [View Range Maps](#)

Noxious Weed: Priority 2B - Aquatic Invasive Species - Non-native Species Global: G5 State: SNA

Predicted Models: 2% Optimal (inductive), 38% Moderate (inductive), 27% Low (inductive)

1		
---	--	--

 V - *Myriophyllum spicatum* (Eurasian Water-milfoil) N2A/AIS

[View in Field Guide](#) [View Predicted Models](#) [View Range Maps](#)

Noxious Weed: Priority 2A - Aquatic Invasive Species - Non-native Species Global: GNR State: SNA

Predicted Models: 26% Low (inductive)

--	--	--

 V - *Nymphaea odorata* (American Water-lily) AIS

[View in Field Guide](#) [View Predicted Models](#) [View Range Maps](#)

Aquatic Invasive Species - Non-native Species Global: G5 State: SNA

Predicted Models: 69% Suitable (introduced range) (deductive)

--	--	--

Noxious Weeds: Priority 1A
 V - *Centaurea solstitialis* (Yellow Starthistle) N1A

[View in Field Guide](#) [View Predicted Models](#) [View Range Maps](#)

Noxious Weed: Priority 1A - Non-native Species Global: GNR State: SNA

Predicted Models: 100% Optimal (inductive)

--	--	--

 V - *Isatis tinctoria* (Dyer's Woad) N1A

[View in Field Guide](#) [View Predicted Models](#) [View Range Maps](#)

Noxious Weed: Priority 1A - Non-native Species Global: GNR State: SNA

Predicted Models: 93% Optimal (inductive), 7% Moderate (inductive)

--	--	--

 V - *Taeniatherum caput-medusae* (Medusahead) N1A

[View in Field Guide](#) [View Predicted Models](#) [View Range Maps](#)

Noxious Weed: Priority 1A - Non-native Species Global: G4G5 State: SNA

Predicted Models: 48% Optimal (inductive), 48% Moderate (inductive), 4% Low (inductive)

--	--	--

 V - *Phragmites australis* ssp. *australis* (European Common Reed) N1A

[View in Field Guide](#) [View Predicted Models](#) [View Range Maps](#)

Noxious Weed: Priority 1A - Non-native Species Global: G5T5 State: SNA

Predicted Models: 11% Optimal (inductive), 79% Moderate (inductive), 10% Low (inductive)

--	--	--

Noxious Weeds: Priority 1B
 V - *Echium vulgare* (Blueweed) N1B

[View in Field Guide](#) [View Predicted Models](#) [View Range Maps](#)

Noxious Weed: Priority 1B - Non-native Species Global: GNR State: SNA

Predicted Models: 100% Optimal (inductive)

1		
---	--	--

 V - *Lythrum salicaria* (Purple Loosestrife) N1B

[View in Field Guide](#) [View Predicted Models](#) [View Range Maps](#)

Noxious Weed: Priority 1B - Non-native Species Global: G5 State: SNA

Predicted Models: 100% Optimal (inductive)

--	--	--

 V - *Polygonum cuspidatum* (Japanese Knotweed) N1B

[View in Field Guide](#) [View Predicted Models](#) [View Range Maps](#)

Noxious Weed: Priority 1B - Non-native Species Global: GNRTRNR State: SNA

Predicted Models: 97% Optimal (inductive), 3% Moderate (inductive)

106		
-----	--	--














 V - *Cytisus scoparius* (Scotch Broom) N1B











[View in Field Guide](#) [View Predicted Models](#) [View Range Maps](#)

Noxious Weed: Priority 1B - Non-native Species Global: GNR State: SNA

Predicted Models: 34% Optimal (inductive), 60% Moderate (inductive), 6% Low (inductive)

--	--	--

<input type="checkbox"/> V - Polygonum x bohemicum (Bohemian Knotweed) N1B		View in Field Guide View Predicted Models View Range Maps
Noxious Weed: Priority 1B - Non-native Species	Global: GNA State: SNA	
Predicted Models:  25% Optimal (inductive),  32% Moderate (inductive),  15% Low (inductive)		
<input type="checkbox"/> V - Chondrilla juncea (Rush Skeletonweed) N1B		View in Field Guide View Predicted Models View Range Maps
Noxious Weed: Priority 1B - Non-native Species	Global: GNR State: SNA	N
Predicted Models:  80% Moderate (inductive),  20% Low (inductive)		
Noxious Weeds: Priority 2A		
<input type="checkbox"/> V - Iris pseudacorus (Yellowflag Iris) N2A/AIS	29 	View in Field Guide View Predicted Models View Range Maps
Noxious Weed: Priority 2A - Aquatic Invasive Species - Non-native Species	Global: GNR State: SNA	N
Predicted Models:  100% Optimal (inductive)		
<input type="checkbox"/> V - Rhamnus cathartica (Common Buckthorn) N2A	192 	View in Field Guide View Predicted Models View Range Maps
Noxious Weed: Priority 2A - Non-native Species	Global: GNR State: SNA	N
Predicted Models:  100% Optimal (inductive)		
<input type="checkbox"/> V - Venenata dubia (Venenata) N2A		View in Field Guide View Predicted Models View Range Maps
Noxious Weed: Priority 2A - Non-native Species	Global: GNR State: SNA	N
Predicted Models:  100% Optimal (inductive)		
<input type="checkbox"/> V - Lepidium latifolium (Perennial Pepperweed) N2A	234 	View in Field Guide View Predicted Models View Range Maps
Noxious Weed: Priority 2A - Non-native Species	Global: GNR State: SNA	N
Predicted Models:  26% Optimal (inductive),  34% Moderate (inductive),  34% Low (inductive)		
<input type="checkbox"/> V - Butomus umbellatus (Flowering-rush) N2A/AIS		View in Field Guide View Predicted Models View Range Maps
Noxious Weed: Priority 2A - Aquatic Invasive Species - Non-native Species	Global: G5 State: SNA	N
Predicted Models:  11% Optimal (inductive),  23% Moderate (inductive),  26% Low (inductive)		
<input type="checkbox"/> V - Hieracium praealtum (Kingdevil Hawkweed) N2A		View in Field Guide View Predicted Models View Range Maps
Noxious Weed: Priority 2A - Non-native Species	Global: GNR State: SNA	N
Predicted Models:  1% Optimal (inductive),  94% Moderate (inductive),  5% Low (inductive)		
<input type="checkbox"/> V - Ranunculus acris (Tall Buttercup) N2A	1 	View in Field Guide View Predicted Models View Range Maps
Noxious Weed: Priority 2A - Non-native Species	Global: G5 State: SNA	N
Predicted Models:  38% Moderate (inductive),  62% Low (inductive)		
<input type="checkbox"/> V - Hieracium aurantiacum (Orange Hawkweed) N2A	17 	View in Field Guide View Predicted Models View Range Maps
Noxious Weed: Priority 2A - Non-native Species	Global: GNR State: SNA	N
Predicted Models:  12% Moderate (inductive),  88% Low (inductive)		
<input type="checkbox"/> V - Senecio jacobaea (Tansy Ragwort) N2A		View in Field Guide View Predicted Models View Range Maps
Noxious Weed: Priority 2A - Non-native Species	Global: GNR State: SNA	N
Predicted Models:  42% Low (inductive)		
<input type="checkbox"/> V - Myriophyllum spicatum (Eurasian Water-milfoil) N2A/AIS		View in Field Guide View Predicted Models View Range Maps
Noxious Weed: Priority 2A - Aquatic Invasive Species - Non-native Species	Global: GNR State: SNA	N
Predicted Models:  26% Low (inductive)		
Noxious Weeds: Priority 2B		
<input type="checkbox"/> V - Tanacetum vulgare (Common Tansy) N2B	1091 	View in Field Guide View Predicted Models View Range Maps
Noxious Weed: Priority 2B - Non-native Species	Global: GNR State: SNA	N
Predicted Models:  100% Optimal (inductive)		
<input type="checkbox"/> V - Tamarix ramosissima (Salt Cedar) N2B	36 	View in Field Guide View Predicted Models View Range Maps
Noxious Weed: Priority 2B - Non-native Species	Global: GNR State: SNA	N
Predicted Models:  34% Optimal (inductive),  59% Moderate (inductive),  7% Low (inductive)		
<input type="checkbox"/> V - Linaria dalmatica (Dalmatian Toadflax) N2B	318 	View in Field Guide View Predicted Models View Range Maps
Noxious Weed: Priority 2B - Non-native Species	Global: G5 State: SNA	N
Predicted Models:  33% Optimal (inductive),  67% Moderate (inductive)		
<input type="checkbox"/> V - Lepidium draba (Whitetop) N2B	11 	View in Field Guide View Predicted Models View Range Maps
Noxious Weed: Priority 2B - Non-native Species	Global: GNR State: SNA	N
Predicted Models:  6% Optimal (inductive),  94% Moderate (inductive)		
<input type="checkbox"/> V - Linaria vulgaris (Yellow Toadflax) N2B	2 	View in Field Guide View Predicted Models View Range Maps
Noxious Weed: Priority 2B - Non-native Species	Global: GNR State: SNA	N
Predicted Models:  3% Optimal (inductive),  34% Moderate (inductive),  49% Low (inductive)		

<input type="checkbox"/> V - Potamogeton crispus (Curly-leaf Pondweed) N2B/AIS		
View in Field Guide View Predicted Models View Range Maps		
Noxious Weed: Priority 2B - Aquatic Invasive Species - Non-native Species Global: G5 State: SNA		
Predicted Models:  2% Optimal (inductive),  38% Moderate (inductive),  27% Low (inductive)		
<input type="checkbox"/> V - Centaurea diffusa (Diffuse Knapweed) N2B		
View in Field Guide View Predicted Models View Range Maps		
Noxious Weed: Priority 2B - Non-native Species Global: GNR State: SNA		
Predicted Models:  89% Moderate (inductive),  11% Low (inductive)		
<input type="checkbox"/> V - Leucanthemum vulgare (Oxeye Daisy) N2B		
View in Field Guide View Predicted Models View Range Maps		
Noxious Weed: Priority 2B - Non-native Species Global: GNR State: SNA		
Predicted Models:  85% Moderate (inductive),  15% Low (inductive)		
<input type="checkbox"/> V - Hypericum perforatum (Common St. John's-wort) N2B		
View in Field Guide View Predicted Models View Range Maps		
Noxious Weed: Priority 2B - Non-native Species Global: GNR State: SNA		
Predicted Models:  74% Moderate (inductive),  26% Low (inductive)		
<input type="checkbox"/> V - Cynoglossum officinale (Common Hound's-tongue) N2B		
View in Field Guide View Predicted Models View Range Maps		
Noxious Weed: Priority 2B - Non-native Species Global: GNR State: SNA		
Predicted Models:  70% Moderate (inductive),  30% Low (inductive)		
<input type="checkbox"/> V - Potentilla recta (Sulphur Cinquefoil) N2B		
View in Field Guide View Predicted Models View Range Maps		
Noxious Weed: Priority 2B - Non-native Species Global: GNR State: SNA		
Predicted Models:  56% Moderate (inductive),  44% Low (inductive)		
<input type="checkbox"/> V - Acroptilon repens (Russian Knapweed) N2B		
View in Field Guide View Predicted Models View Range Maps		
Noxious Weed: Priority 2B - Non-native Species Global: GNR State: SNA		
Predicted Models:  25% Moderate (inductive),  61% Low (inductive)		
<input type="checkbox"/> V - Centaurea stoebe (Spotted Knapweed) N2B		
View in Field Guide View Predicted Models View Range Maps		
Noxious Weed: Priority 2B - Non-native Species Global: GNR State: SNA		
Predicted Models:  22% Moderate (inductive),  63% Low (inductive)		
<input type="checkbox"/> V - Euphorbia virgata (Leafy Spurge) N2B		
View in Field Guide View Predicted Models View Range Maps		
Noxious Weed: Priority 2B - Non-native Species Global: GNR State: SNA		
Predicted Models:  3% Moderate (inductive),  97% Low (inductive)		
<input type="checkbox"/> V - Cirsium arvense (Canada Thistle) N2B		
View in Field Guide View Predicted Models View Range Maps		
Noxious Weed: Priority 2B - Non-native Species Global: G5 State: SNA		
Predicted Models:  3% Moderate (inductive),  80% Low (inductive)		
<input type="checkbox"/> V - Convolvulus arvensis (Field Bindweed) N2B		
View in Field Guide View Predicted Models View Range Maps		
Noxious Weed: Priority 2B - Non-native Species Global: GNR State: SNA		
Predicted Models:  88% Low (inductive)		
<input type="checkbox"/> V - Berteroa incana (Hoary False-alyssum) N2B		
View in Field Guide View Predicted Models View Range Maps		
Noxious Weed: Priority 2B - Non-native Species Global: GNR State: SNA		
Predicted Models:  67% Low (inductive)		
Regulated Weeds: Priority 3		
<input type="checkbox"/> V - Elaeagnus angustifolia (Russian Olive) R3		
View in Field Guide View Predicted Models View Range Maps		
Regulated Weed: Priority 3 - Non-native Species Global: GNR State: SNA		
Predicted Models:  13% Optimal (inductive),  19% Moderate (inductive),  67% Low (inductive)		
<input type="checkbox"/> V - Bromus tectorum (Cheatgrass) R3		
View in Field Guide View Predicted Models View Range Maps		
Regulated Weed: Priority 3 - Non-native Species Global: GNR State: SNA		
Predicted Models:  69% Moderate (inductive),  31% Low (inductive)		
Forest Pests		
<input type="checkbox"/> I - Leucoma salicis (Satin Moth) PESTF		
View in Field Guide		
Forest Pest - Native Species Global: GNR State: SNR		
Biocontrol Species		
<input type="checkbox"/> I - Cyphocleonus achates (Knapweed Root Weevil) BIOCNTRL		
View in Field Guide View Predicted Models View Range Maps		
Biocontrol Species - Non-native Species Global: GNR State: SNA		
Predicted Models:  100% Optimal (inductive)		
<input type="checkbox"/> I - Oberea erythrocephala (Red-headed Leafy Spurge Stem Borer) BIOCNTRL		
View in Field Guide View Predicted Models View Range Maps		
Biocontrol Species - Non-native Species Global: GNR State: SNA		
Predicted Models:  100% Optimal (inductive)		

[View in Field Guide](#) [View Predicted Models](#) [View Range Maps](#)Biocontrol Species - Non-native Species Global: **GNR** State: **SNA**Predicted Models:  54% Optimal (inductive),  46% Moderate (inductive)[View in Field Guide](#) [View Predicted Models](#) [View Range Maps](#)Biocontrol Species - Non-native Species Global: **GNR** State: **SNA**Predicted Models:  21% Optimal (inductive),  75% Moderate (inductive),  4% Low (inductive)[View in Field Guide](#) [View Predicted Models](#) [View Range Maps](#)Biocontrol Species - Non-native Species Global: **GNR** State: **SNA**Predicted Models:  100% Moderate (inductive)[View in Field Guide](#) [View Predicted Models](#) [View Range Maps](#)Biocontrol Species - Non-native Species Global: **GNR** State: **SNA**Predicted Models:  77% Moderate (inductive),  23% Low (inductive)[View in Field Guide](#)Biocontrol Species - Non-native Species Global: **G5** State: **SNA**

Introduction to Montana Natural Heritage Program



PO Box 201800 • 1201 11th Avenue • Helena, MT 59620-1800 • fax 406.444.0266 • phone 406.444.3989 • mtnhp.org

INTRODUCTION

The Montana Natural Heritage Program (MTNHP) is Montana's source for reliable and objective information on Montana's native species and habitats, emphasizing those of conservation concern. MTNHP was created by the Montana legislature in 1983 as part of the Natural Resource Information System (NRIS) at the Montana State Library (MSL). MTNHP is "a program of information acquisition, storage, and retrieval for data relating to the flora, fauna, and biological community types of Montana" (MCA 90-15-102). MTNHP's activities are guided by statute as well as through ongoing interaction with, and feedback from, principal data source agencies such as Montana Fish, Wildlife, and Parks, the Montana Department of Environmental Quality, the Montana Department of Natural Resources and Conservation, the Montana University System, the US Forest Service, and the US Bureau of Land Management. Since the first staff was hired in 1985, the Program has logged a long record of success, and developed into a highly respected, service-oriented program. MTNHP is widely recognized as one of the most advanced and effective of over 60 natural heritage programs that are distributed across North America.

VISION

Our vision is that public agencies, the private sector, the education sector, and the general public will trust and rely upon MTNHP as the source for information and expertise on Montana's species and habitats, especially those of conservation concern. We strive to provide easy access to our information to allow users to save time and money, speed environmental reviews, and make informed decisions.

CORE VALUES

- We endeavor to be a single statewide source of accurate and up-to-date information on Montana's plants, animals, and aquatic and terrestrial biological communities.
- We actively listen to our data users and work responsively to meet their information and training needs.
- We strive to provide neutral, trusted, timely, and equitable service to all of our information users.
- We make every effort to be transparent to our data users in setting work priorities and providing data products.

CONFIDENTIALITY

All information requests made to the Montana Natural Heritage Program are considered library records and are protected from disclosure by the Montana Library Records Confidentiality Act (MCA 22-1-11).

INFORMATION MANAGED

Information managed at the Montana Natural Heritage Program is botanical, zoological, and ecological information that describes the distribution (e.g., observations, structured surveys, range polygons, predicted habitat suitability models), conservation status (e.g., global and state conservation status ranks, including threats), and other supporting information (e.g., accounts and references) on the biology and ecology of species and biological communities.

Data Use Terms and Conditions

- Montana Natural Heritage Program (MTNHP) products and services are based on biological data and the objective interpretation of those data by professional scientists. MTNHP does not advocate any particular philosophy of natural resource protection, management, development, or public policy.
- MTNHP has no natural resource management or regulatory authority. Products, statements, and services from MTNHP are intended to inform parties as to the state of scientific knowledge about certain natural resources, and to further develop that knowledge. The information is not intended as natural resource management guidelines or prescriptions or a determination of environmental impacts. MTNHP recommends consultation with appropriate state, federal, and tribal resource management agencies and authorities in the area where your project is located.
- Information on the status and spatial distribution of biological resources produced by MTNHP are intended to inform parties of the state-wide status, known occurrence, or the likelihood of the presence of those resources. **These products are not intended to substitute for field-collected data, nor are they intended to be the sole basis for natural resource management decisions.**
- MTNHP does not portray its data as exhaustive or comprehensive inventories of rare species or biological communities. **Field verification of the absence or presence of sensitive species and biological communities will always be an important obligation of users of our data.**
- MTNHP responds equally to all requests for products and services, regardless of the purpose or identity of the requester.
- Because MTNHP constantly updates and revises its databases with new data and information, products will become outdated over time. Interested parties are encouraged to obtain the most current information possible from MTNHP, rather than using older products. We add, review, update, and delete records on a daily basis. Consequently, we strongly advise that you update your MTNHP data sets at a minimum of every four months for most applications of our information.
- MTNHP data require a certain degree of biological expertise for proper analysis, interpretation, and application. Our staff is available to advise you on questions regarding the interpretation or appropriate use of the data that we provide. See [Contact Information for MTNHP Staff](#)
- The information provided to you by MTNHP may include sensitive data that if publicly released might jeopardize the welfare of threatened, endangered, or sensitive species or biological communities. This information is intended for distribution or use only within your department, agency, or business. Subcontractors may have access to the data during the course of any given project, but should not be given a copy for their use on subsequent, unrelated work.
- MTNHP data are made freely available. Duplication of hard-copy or digital MTNHP products with the intent to sell is prohibited without written consent by MTNHP. Should you be asked by individuals outside your organization for the type of data that we provide, please refer them to MTNHP.
- MTNHP and appropriate staff members should be appropriately acknowledged as an information source in any third-party product involving MTNHP data, reports, papers, publications, or in maps that incorporate MTNHP graphic elements.
- Sources of our data include museum specimens, published and unpublished scientific literature, field surveys by state and federal agencies and private contractors, and reports from knowledgeable individuals. MTNHP actively solicits and encourages additions, corrections and updates, new observations or collections, and comments on any of the data we provide.
- MTNHP staff and contractors do not enter or cross privately-owned lands without express permission from the landowner. However, the program cannot guarantee that information provided to us by others was obtained under adherence to this policy.

Suggested Contacts for Natural Resource Management Agencies

As required by Montana statute (MCA 90-15), the Montana Natural Heritage Program works with state, federal, tribal, nongovernmental organizations, and private partners to ensure that the latest animal and plant distribution and status information is incorporated into our databases so that it can be used to inform a variety of permitting and planning processes and management decisions. We encourage you to contact state, federal, and tribal resource management agencies in the area where your project is located and review the permitting overviews by the [Montana Department of Environmental Quality](#), the [Montana Department of Natural Resources and Conservation](#) and the [Index of Environmental Permits for Montana](#) for guidelines relevant to your efforts. In particular, we encourage you to contact the Montana Department of Fish, Wildlife, and Parks for the latest data and management information regarding hunted and high-profile management species and to use the U.S. Fish and Wildlife Service's [Information Planning and Consultation \(IPAC\) website regarding U.S. Endangered Species Act listed Threatened, Endangered, or Candidate species](#).

For your convenience, we have compiled a list of relevant agency contacts and links below:

Montana Fish, Wildlife, and Parks

Fish Species	Zachary Shattuck zshattuck@mt.gov (406) 444-1231 or Eric Roberts eroberts@mt.gov (406) 444-5334
American Bison Black-footed Ferret Black-tailed Prairie Dog Bald Eagle Golden Eagle Common Loon Least Tern Piping Plover Whooping Crane	Kristian Smucker KSmucker@mt.gov (406) 444-5209
Grizzly Bear Greater Sage Grouse Trumpeter Swan Big Game Upland Game Birds Furbearers	Brian Wakeling brian.wakeling@mt.gov (406) 444-3940
Managed Terrestrial Game Data	Cara Whalen– MFWP Data Analyst cara.whalen@mt.gov (406) 444-3759
Fisheries Data and Nongame Animal Data	Ryan Alger – MFWP Data Analyst ryan.alger@mt.gov (406) 444-5365
Wildlife and Fisheries Scientific Collector's Permits	https://fwp.mt.gov/buyandapply/commercialwildlifeandscientificpermits/scientific Kristina Smucker for Wildlife ksmucker@mt.gov (406) 444-5209 Dave Schmetterling for Fisheries dschmetterling@mt.gov (406) 542-5514
Fish and Wildlife Recommendations for Subdivision Development	Charlie Sperry csperry@mt.gov (406) 444-3888 See https://fwp.mt.gov/conservation/living-with-wildlife/subdivision-recommendations
Regional Contacts 	Region 1 (Kalispell) (406) 752-5501 fwprg12@mt.gov Region 2 (Missoula) (406) 542-5500 fwprg22@mt.gov Region 3 (Bozeman) (406) 577-7900 fwprg3@mt.gov Region 4 (Great Falls) (406) 454-5840 fwprg42@mt.gov Region 5 (Billings) (406) 247-2940 fwprg52@mt.gov Region 6 (Glasgow) (406) 228-3700 fwprg62@mt.gov Region 7 (Miles City) (406) 234-0900 fwprg72@mt.gov

Montana Department of Agriculture

General Contact Information: <https://agr.mt.gov/About/Office-Locations/Office-Locations-and-Field-Offices>

Noxious Weeds: <https://agr.mt.gov/Noxious-Weeds>

Montana Department of Environmental Quality

Permitting and Operator Assistance for all Environmental Permits: <https://deq.mt.gov/Permitting>

Montana Department of Natural Resources and Conservation

Overview of, and contacts for, licenses and permits for state lands, water, and forested lands:

<https://dnrc.mt.gov/Permits-Services>

Stream Permitting (310 permits) and an overview of various water and stream related permits (e.g., Stream Protection Act 124, Federal Clean Water Act 404, Federal Rivers and Harbors Act Section 10, Short-term Water Quality Standard for Turbidity 318 Authorization, etc.).

<https://dnrc.mt.gov/Licenses-and-Permits/Stream-Permitting>

Wildfire Resources: <https://dnrc.mt.gov/Forestry/Wildfire>

Bureau of Land Management



Billings	(406) 896-5013
Butte	(406) 533-7600
Dillon	(406) 683-8000
Glasgow	(406) 228-3750
Havre	(406) 262-2820
Lewistown	(406) 538-1900
Malta	(406) 654-5100
Miles City	(406) 233-2800
Missoula	(406) 329-3914

United States Army Corps of Engineers

Montana Regulatory Office for federal permits related to construction in water and wetlands

<https://www.nwo.usace.army.mil/Missions/Regulatory-Program/Montana/> (406) 441-1375

United States Environmental Protection Agency

Environmental information, notices, permitting, and contacts <https://www.epa.gov/mt>

Gateway to state resource locators <https://www.envcap.org/srl/index.php>

United States Fish and Wildlife Service

Information Planning and Conservation (IPAC) website: <https://ipac.ecosphere.fws.gov>


Montana Ecological Services Field Office: <https://www.fws.gov/office/montana-ecological-services> (406) 449-5225

United States Forest Service

Regional Office – Missoula, Montana Contacts

Wildlife Program Leader	Tammy Fletcher	tammy.fletcher2@usda.gov	(406) 329-3086
Wildlife Ecologist	Cara Staab	cara.staab@usda.gov	(406) 329-3677
Aquatic Ecologist	Justin Jimenez	justin.jimenez@usda.gov	(435) 370-6830
TES Program	Lydia Allen	lydia.allen@usda.gov	(406) 329-3558
Interagency Grizzly Bear Coordinator	Scott Jackson	scott.jackson@usda.gov	(406) 329-3664
Regional Botanist	Amanda Hendrix	amanda.hendrix@usda.gov	(651) 447-3016
Regional Vegetation Ecologist	Mary Manning	mary.manning@usda.gov	(406) 329-3304
Invasive Species Program Manager	Michelle Cox	micelle.cox2@usda.gov	(406) 329-3669

Tribal Nations



- [Assiniboine & Gros Ventre Tribes – Fort Belknap Reservation](#)
- [Assiniboine & Sioux Tribes – Fort Peck Reservation](#)
- [Blackfeet Tribe - Blackfeet Reservation](#)
- [Chippewa Creek Tribe - Rocky Boy's Reservation](#)
- [Crow Tribe – Crow Reservation](#)
- [Little Shell Chippewa Tribe](#)
- [Northern Cheyenne Tribe – Northern Cheyenne Reservation](#)
- [Salish & Kootenai Tribes - Flathead Reservation](#)

Natural Heritage Programs and Conservation Data Centers in Surrounding States and Provinces

[Alberta Conservation Information Management System](#)

[British Columbia Conservation Data Centre](#)

[Idaho Natural Heritage Program](#)

[North Dakota Natural Heritage Program](#)

[Saskatchewan Conservation Data Centre](#)

[South Dakota Natural Heritage Program](#)

[Wyoming Natural Diversity Database](#)

Invasive Species Management Contacts and Information

[Aquatic Invasive Species](#)

[Montana Fish, Wildlife, and Parks Aquatic Invasive Species staff](#)

[Montana Department of Natural Resources and Conservation's Aquatic Invasive Species Grant Program](#)

[Montana Invasive Species Council \(MISC\)](#)

[Upper Columbia Conservation Commission \(UC3\)](#)

[Noxious Weeds](#)

[Montana Weed Control Association Contacts Webpage](#)

[Montana Biological Weed Control Coordination Project](#)

[Montana Department of Agriculture - Noxious Weeds](#)

[Montana Weed Control Association](#)

[Montana Fish, Wildlife, and Parks - Noxious Weeds](#)

[Montana State University Integrated Pest Management Extension](#)

[Integrated Noxious Weed Management after Wildfires](#)

[Fire Management and Invasive Plants](#)

Introduction to Native Species

Within the report area you have requested, separate summaries are provided for: (1) Species Occurrences (SO) for plant and animal Species of Concern, Special Status Species (SSS), Important Animal Habitat (IAH) and some Potential Plant Species of Concern; (2) other observed non Species of Concern or Species of Concern without suitable documentation to create Species Occurrence polygons; and (3) other non-documented species that are potentially present based on their range, predicted suitable habitat model output, or presence of associated habitats. Each of these summaries provides the following information when present for a species: (1) the number of [Species Occurrences](#) and associated delineation criteria for construction of these polygons that have long been used for considerations of documented Species of Concern in environmental reviews; (2) the number of observations of each species; (3) the geographic range polygons for each species that the report area overlaps; (4) predicted relative habitat suitability classes that are present if a predicted suitable habitat model has been created; (5) the percent of the report area that is mapped as commonly associated or occasionally associated habitat as listed for each species in the [Montana Field Guide](#); and (6) a variety of conservation status ranks and links to species accounts in the [Montana Field Guide](#). Details on each of these information categories are included under relevant section headers below or are defined on our [Species Status Codes](#) page. In presenting this information, the Montana Natural Heritage Program (MTNHP) is working towards assisting the user with rapidly determining what species have been documented and what species are potentially present in the report area. We remind users that this information is likely incomplete as surveys to document native and introduced species are lacking in many areas of the state, information on introduced species has only been tracked relatively recently, the MTNHP's staff and resources are restricted by budgets, and information is constantly being added and updated in our databases. **Thus, field verification by professional biologists of the absence or presence of species and biological communities will always be an important obligation of users of our data.**

If you are aware of observation datasets that the MTNHP is missing, please report them to the Program Botanist apipp@mt.gov or Senior Zoologist dbachen@mt.gov If you have animal or plant observations that you would like to contribute, you can also submit them via Excel spreadsheets, geodatabases, iNaturalist, or a Survey123 form. Various methods of data submission are reviewed in this playlist of videos:

<https://www.youtube.com/playlist?list=PLRaydtZpHu2qOHPoSPq9cnM9uXGmEXACx>

Observations

The MTNHP manages information on several million animal and plant observations that have been reported by professional biologists and private citizens from across Montana. The majority of these observations are submitted in digital format from standardized databases associated with research or monitoring efforts and spreadsheets of incidental observations submitted by professional biologists and amateur naturalists. At a minimum, accepted observation records must contain a credible species identification (i.e. appropriate geographic range, date, and habitat and, if species are difficult to identify, a photograph and/or notes on key identifying features), a date or date range, observer name, locational information (ideally with latitude and longitude in decimal degrees), notes on numbers observed, and species behavior or habitat use (e.g., is the observation likely associated with reproduction). Bird records are also required to have information associated with date-appropriate breeding or overwintering status of the species observed. MTNHP reviews observation records to ensure that they are mapped correctly, occur within date ranges when the species is known to be present or detectable, occur within the known seasonal geographic range of the species, and occur in appropriate habitats. MTNHP also assigns each record a locational uncertainty value in meters to indicate the spatial precision associated with the record's mapped coordinates. Only records with locational uncertainty values of 10,000 meters or less are included in environmental summary reports and number summaries are only provided for records with locational uncertainty values of 1,000 meters or less.

Species Occurrences

The MTNHP evaluates plant and animal observation records for species of higher conservation concern to determine whether they are worthy of inclusion in the [Species Occurrence](#) (SO) layer for use in environmental reviews; observations not worthy of inclusion in this layer include long distance dispersal events, migrants observed away from key migratory stopover habitats, and winter observations. An SO is a polygon depicting what is known about a species occupancy from direct observation with a defined level of locational uncertainty and any inference that can be made about adjacent habitat use from the latest peer-reviewed science. If an observation can be associated with a map feature that can be tracked (e.g., a wetland boundary for a wetland associated plant) then this polygon feature is used to represent the SO. Areas that can be inferred as probable occupied habitat based on direct observation of a species location and what is known about the foraging area or home range size of the species may be incorporated into the SO. Species Occurrences generally belong to one of the following categories:

Plant Species Occurrences

A documented location of a specimen collection or observed plant population. In some instances, adjacent, spatially separated clusters are considered subpopulations and are grouped as one occurrence (e.g., the subpopulations occur in ecologically similar habitats, and their spatial proximity likely allows them to interbreed). Tabular information for multiple observations at the same SO location is generally linked to a single polygon. Plant SO's are only created for Species of Concern and Potential Species of Concern.

Animal Species Occurrences


The location of a verified observation or specimen record typically known or assumed to represent a breeding population or a portion of a breeding population. Animal SO's are generally: (1) buffers of terrestrial point observations based on documented species' home range sizes; (2) buffers of stream segments to encompass occupied streams and immediate adjacent riparian habitats; (3) polygonal features encompassing known or likely breeding populations (e.g., a wetland for some amphibians or a forested portion of a mountain range for some wide-ranging carnivores); or (4) combinations of the above. Tabular information for multiple observations at the same SO location is generally linked to a single polygon. Species Occurrence polygons may encompass some unsuitable habitat in some instances in order to avoid heavy data processing associated with clipping out habitats that are readily assessed as unsuitable by the data user (e.g., a point buffer of a terrestrial species may overlap into a portion of a lake that is obviously inappropriate habitat for the species). Animal SO's are only created for Species of Concern and Special Status Species (e.g., Bald Eagle).

Other Occurrence Polygons

These include significant biological features not included in the above categories, such as Important Animal Habitats like bird rookeries and bat roosts, and peatlands or other wetland and riparian communities that support diverse plant and animal communities.

Geographic Range Polygons

Geographic range polygons are still under development for most plant and invertebrate species. Native year-round, summer, winter, migratory and historic geographic range polygons as well as polygons for introduced



populations have been defined for most vertebrate animal species for which there are enough observations, surveys, and knowledge of appropriate seasonal habitat use to define them (see examples to left). These native or introduced range polygons bound the extent of known or likely occupied habitats for non-migratory and relative sedentary species and the regular extent of known or likely occupied habitats for migratory and long-distance dispersing species; polygons may include unsuitable intervening habitats. For most species, a single polygon can represent the year-round or seasonal range, but breeding ranges of some colonial nesting water birds and some introduced species are represented more patchily when supported by data. Some ranges are mapped more broadly than actual distributions in order to be visible on statewide maps (e.g., fish).

Predicted Suitable Habitat Models

Predicted habitat suitability models have been created for plant and animal Species of Concern and are undergoing development for non-Species of Concern. For species for which models have been completed, the environmental summary report includes simple rule-based associations with streams for aquatic species and seasonal habitats for game species as well as mathematically complex Maximum Entropy models (Phillips et al. 2006, Ecological Modeling 190:231-259) constructed from a variety of statewide biotic and abiotic layers and presence only data for individual species for most terrestrial species. For the Maximum Entropy models, we reclassified 90 x 90-meter continuous model output into suitability classes (unsuitable, low, moderate, and optimal) then aggregated that into the one square mile hexagons used in the environmental summary report; this is the finest spatial scale we suggest using this information in management decisions and survey planning. Full model write ups for individual species that discuss model goals, inputs, outputs, and evaluation in much greater detail are posted on the MTNHP's [Predicted Suitable Habitat Models](#) webpage. Evaluations of predictive accuracy and specific limitations are included with the metadata for models of individual species.

Model outputs should not be used in place of on-the-ground surveys for species. Instead model outputs should be used in conjunction with habitat evaluations to determine the need for on-the-ground surveys for species. We suggest that the percentage of predicted optimal and moderate suitable habitat within the report area be used in conjunction with geographic range polygons and the percentage of commonly associated habitats to generate lists of potential species that may occupy broader landscapes for the purposes of landscape-level planning.

Associated Habitats

Within the boundary of the intersected hexagons, we provide the approximate percentage of commonly or occasionally associated habitat for vertebrate animal species that regularly breed, overwinter, or migrate through the state; a detailed list of commonly and occasionally associated habitats is provided in individual species accounts in the [Montana Field Guide](#). We assigned common or occasional use of each of the ecological

systems mapped in Montana by: (1) using personal knowledge and reviewing literature that summarizes the breeding, overwintering, or migratory habitat requirements of each species; (2) evaluating structural characteristics and distribution of each ecological system relative to the species' range and habitat requirements; (3) examining the observation records for each species in the state-wide point observation database associated with each ecological system; and (4) calculating the percentage of observations associated with each ecological system relative to the percent of Montana covered by each ecological system to get a measure of numbers of observations versus availability of habitat. Species that breed in Montana were only evaluated for breeding habitat use, species that only overwinter in Montana were only evaluated for overwintering habitat use, and species that only migrate through Montana were only evaluated for migratory habitat use. In general, species were listed as associated with an ecological system if structural characteristics of used habitat documented in the literature were present in the ecological system or large numbers of point observations were associated with the ecological system. However, species were not listed as associated with an ecological system if there was no support in the literature for use of structural characteristics in an ecological system, even if point observations were associated with that system. Common versus occasional association with an ecological system was assigned based on the degree to which the structural characteristics of an ecological system matched the preferred structural habitat characteristics for each species as represented in the scientific literature. The percentage of observations associated with each ecological system relative to the percent of Montana covered by each ecological system was also used to guide assignment of common versus occasional association.

We suggest that the percentage of commonly associated habitat within the report area be used in conjunction with geographic range polygons and the percentage of predicted optimal and moderate suitable habitat from predictive models to generate lists of potential species that may occupy broader landscapes for the purposes of landscape-level planning. Users of this information should be aware that land cover mapping accuracy is particularly problematic when the systems occur as small patches or where the land cover types have been altered over the past decade. Thus, particular caution should be used when using the associations in assessments of smaller areas (e.g., evaluations of public land survey sections).

Introduction to Land Cover

Land Use/Land Cover is one of 15 [Montana Spatial Data Infrastructure](#) framework layers considered vital for making statewide maps of Montana and understanding its geography. The layer records all Montana natural vegetation, land cover and land use, classified from satellite and aerial imagery, mapped at a scale of 1:100,000, and interpreted with supporting ground-level data. The baseline map is adapted from the Northwest ReGAP (NWGAP) project land cover classification, which used 30m resolution multi-spectral Landsat imagery acquired between 1999 and 2001. Vegetation classes were drawn from the Ecological System Classification developed by NatureServe (Comer et al. 2003). The land cover classes were developed by Anderson et al. (1976). The NWGAP effort encompasses 12 map zones. Montana overlaps seven of these zones. The two NWGAP teams responsible for the initial land cover mapping effort in Montana were Sanborn and NWGAP at the University of Idaho. Both Sanborn and NWGAP employed a similar modeling approach in which Classification and Regression Tree (CART) models were applied to Landsat ETM+ scenes. The Spatial Analysis Lab within the Montana Natural Heritage Program was responsible for developing a seamless Montana land cover map with a consistent statewide legend from these two separate products. Additionally, the Montana land cover layer incorporates several other land cover and land use products (e.g., MSDI Structures and Transportation themes and the Montana Department of Revenue Final Land Unit classification) and reclassifications based on plot-level data and the latest NAIP imagery to improve accuracy and enhance the usability of the theme. Updates are done as partner support and funding allow, or when other MSDI datasets can be incorporated. Recent updates include fire perimeters and agricultural land use (annually), energy developments such as wind, oil and gas installations (2014), roads, structures and other impervious surfaces (various years); and local updates/improvements to specific ecological systems (e.g., central Montana grassland and sagebrush ecosystems). Current and previous versions of the Land Use/Land Cover layer with full metadata are available for download from the Montana State Library's [GIS Data List](#). More information on the land cover layer is available at: https://msl.mt.gov/geoinfo/msdi/land_use_land_cover/

Within the report area you have requested, land cover is summarized by acres of Level 1, Level 2, and Level 3 Ecological Systems.

Literature Cited

Anderson, J.R. E.E. Hardy, J.T. Roach, and R.E. Witmer. 1976. A land use and land cover classification system for use with remote sensor data. U.S. Geological Survey Professional Paper 964.

Comer, P., D. Faber-Langendoen, R. Evans, S. Gawler, C. Josse, G. Kittel, S. Menard, M. Pyne, M. Reid, K. Schulz, K. Snow, and J. Teague. 2003. Ecological systems of the United States: A working classification of U.S. terrestrial systems. NatureServe, Arlington, VA.

Introduction to Wetland and Riparian

Within the report area you have requested, wetland and riparian mapping is summarized by acres of each classification present. Summaries are only provided for modern MTNHP wetland and riparian mapping and not for outdated (NWI Legacy) or incomplete (NWI Scalable) mapping efforts; [described here](#). MTNHP has made all three of these datasets and associated metadata available for separate download on the Montana [Wetland and Riparian Framework](#) web page.

Wetland and Riparian mapping is one of 15 [Montana Spatial Data Infrastructure](#) framework layers considered vital for making statewide maps of Montana and understanding its geography. The wetland and riparian framework layer consists of spatial data representing the extent, type, and approximate location of wetlands, riparian areas, and deep water habitats in Montana.

Wetland and riparian mapping is completed through photointerpretation of 1-m resolution color infrared aerial imagery acquired from 2005 or later. A coding convention using letters and numbers is assigned to each mapped wetland. These letters and numbers describe the broad landscape context of the wetland, its vegetation type, its water regime, and the kind of alterations that may have occurred. Ancillary data layers such as topographic maps, digital elevation models, soils data, and other aerial imagery sources are also used to improve mapping accuracy. Wetland mapping follows the federal Wetland Mapping Standard and classifies wetlands according to the Cowardin classification system of the National Wetlands Inventory (NWI) (Cowardin et al. 1979, FGDC Wetlands Subcommittee 2013). Federal, State, and local regulatory agencies with jurisdiction over wetlands may define and describe wetlands differently than the NWI. Similar coding, based on U.S. Fish and Wildlife Service conventions, is applied to riparian areas (U.S. Fish and Wildlife Service 2009). These are mapped areas where vegetation composition and growth is influenced by nearby water bodies, but where soils, plant communities, and hydrology do not display true wetland characteristics. **These data are intended for use at a scale of 1:12,000 or smaller. Mapped wetland and riparian areas do not represent precise boundaries and digital wetland data cannot substitute for an on-site determination of jurisdictional wetlands.**

See detailed overviews, with examples, of both wetland and riparian classification systems and associated codes as a [storymap](#) and companion [guide](#)

Literature Cited

Cowardin, L.M., V. Carter, F.C. Golet, and E.T. LaRoe. 1979. Classification of wetlands and deepwater habitats of the United States. U.S. Fish and Wildlife Service, FWS/OBS-79/31. Washington, D.C. 103pp.

Federal Geographic Data Committee. 2013. Classification of wetlands and deepwater habitats of the United States. FGDC-STD-004-2013. Second Edition. Wetlands Subcommittee, Federal Geographic Data Committee and U.S. Fish and Wildlife Service, Washington, D.C.

U.S. Fish and Wildlife Services. 2009. A system for mapping riparian areas in the western United States. Division of Habitat and Resource Conservation, Branch of Resource and Mapping Support, Arlington, Virginia.

Introduction to Land Management

Within the report area you have requested, land management information is summarized by acres of federal, state, and local government lands, tribal reservation boundaries, private conservation lands, and federal, state, local, and private conservation easements. Acreage for “Owned”, “Tribal”, or “Easement” categories represents non-overlapping areas that may be totaled. However, “Other Boundaries” represents managed areas such as National Forest boundaries containing private inholdings and other mixed ownership which may cause boundaries to overlap (e.g. a wilderness area within a forest). Therefore, acreages may not total in a straight-forward manner.

Because information on land stewardship is critical to effective land management, the Montana Natural Heritage Program (MTNHP) began compiling ownership and management data in 1997. The goal of the Montana Land Management Database is to manage a single, statewide digital data set that incorporates information from both public and private entities. The database assembles information on public lands, private conservation lands, and conservation easements held by state and federal agencies and land trusts and is updated on a regular basis. Since 2011, the Information Management group in the Montana State Library’s Digital Library Division has led the Montana Land Management Database in partnership with the MTNHP.

Public and private conservation land polygons are attributed with the name of the entity that owns it. The data are derived from the statewide [Montana Cadastral Parcel layer](#). Conservation easement data shows land parcels on which a public agency or qualified land trust has placed a conservation easement in cooperation with the landowner. The dataset contains no information about ownership or status of the mineral estate. For questions about the dataset or to report errors, please contact the Montana Natural Heritage Program at (406) 444-5363 or mtnhp@mt.gov. You can download various components of the Land Management Database and view associated metadata at the Montana State Library’s [GIS Data List](#) at the following links:

[Public Lands](#)

[Conservation Easements](#)

[Private Conservation Lands](#)

[Managed Areas](#)

Map features in the Montana Land Management Database or summaries provided in this report are not intended as a legal depiction of public or private surface land ownership boundaries and should not be used in place of a survey conducted by a licensed land surveyor. Similarly, map features do not imply public access to any lands. The Montana Natural Heritage Program makes no representations or warranties whatsoever with respect to the accuracy or completeness of this data and assumes no responsibility for the suitability of the data for a particular purpose. The Montana Natural Heritage Program will not be liable for any damages incurred as a result of errors displayed here. Consumers of this information should review or consult the primary data and information sources to ascertain the viability of the information for their purposes.

Introduction to Invasive and Pest Species

Within the report area you have requested, separate summaries are provided for: Aquatic Invasive Species, Noxious Weeds, Agricultural Pests, Forest Pests, and Biocontrol species that have been documented or potentially occur there based on the predicted suitability of habitat. Definitions for each of these invasive and pest species categories can be found on our [Species Status Codes](#) page.

Each of these summaries provides the following information when present for a species: (1) the number of observations of each species; (2) the geographic range polygons for each species, if developed, that the report area overlaps; (3) predicted relative habitat suitability classes that are present if a predicted suitable habitat model has been created; (4) the percent of the report area that is mapped as commonly associated or occasionally associated habitat as listed for each species in the [Montana Field Guide](#); and (5) links to species accounts in the [Montana Field Guide](#). Details on each of these information categories are included under relevant section headers under the Introduction to Native Species above or are defined on our [Species Status Codes](#) page. In presenting this information, the Montana Natural Heritage Program (MTNHP) is working towards assisting the user with rapidly determining what invasive and pest species have been documented and what species are potentially present in the report area. We remind users that this information is likely incomplete as surveys to document introduced species are lacking in many areas of the state, information on introduced species has only been tracked relatively recently, the MTNHP's staff and resources are limited, and information is constantly being added and updated in our databases. **Thus, field verification by professional biologists of the absence or presence of species will always be an important obligation of users of our data.**

If you are aware of observation or survey datasets for invasive or pest species that the MTNHP is missing, please report them to the Program Coordinator bmaxell@mt.gov Program Botanist apipp@mt.gov or Senior Zoologist dbachen@mt.gov If you have animal or plant observations that you would like to contribute, you can also submit them via Excel spreadsheets, geodatabases, iNaturalist, or a Survey123 form. Various methods of data submission are reviewed in this playlist of videos:

<https://www.youtube.com/playlist?list=PLRaydtZpHu2qOHPoSPq9cnM9uXGmEXACx>

Additional Information Resources

[MTNHP Staff Contact Information](#)

[Montana Field Guide](#)

[MTNHP Species of Concern Report - Animals and Plants](#)

[MTNHP Species Status Codes - Explanation](#)

[MTNHP Predicted Suitable Habitat Models](#) (for select Animals and Plants)

[MTNHP Request Information page](#)

[Montana Cadastral](#)

[Montana Code Annotated](#)

[Montana Fisheries Information System](#)

[Montana Fish, Wildlife, and Parks Subdivision Recommendations](#)

[Montana GIS Data Layers](#)

[Montana GIS Data Bundler](#)

[Montana Greater Sage-Grouse Project Submittal Site](#)

[Montana Ground Water Information Center](#)

[Montana Index of Environmental Permits, 21st Edition \(2018\)](#)

[Montana Environmental Policy Act \(MEPA\)](#)

[Montana Environmental Policy Act Analysis Resource List](#)

[Laws, Treaties, Regulations, and Agreements on Animals and Plants](#)

[Montana Spatial Data Infrastructure Layers](#)

[Montana State Historic Preservation Office Review and Compliance](#)

[Montana Stream Permitting: a guide for conservation district supervisors and others](#)

[Montana Water Information System](#)

[Montana Web Map Services](#)

[National Environmental Policy Act](#)

[Penalties for Misuse of Fish and Wildlife Location Data](#) (MCA 87-6-222)

[U.S. Fish and Wildlife Service Information for Planning and Consultation](#) (Section 7 Consultation)

[Web Soil Survey Tool](#)